arXiv:1304.3334v1 [quant-ph] 11 Apr 2013

Geometric quantum discord with Bures distance

D. Spehner[1, 2, *] and M. Orszag[3]

[1] *Université Grenoble 1 and CNRS, Institut Fourier UMR5582,*
B.P. 74, 38402 Saint Martin d'Hères, France
[2] *Université Grenoble 1 and CNRS, Laboratoire de Physique et Modélisation des Milieux Condensés UMR5493,*
B.P. 166, 38042 Grenoble, France
[3] *Pontificia Universidad Católica, Facultad de física,*
Casilla 306, Santiago 22, Chile

(Dated: April 12, 2013)

We define a new measure of quantum correlations in bipartite quantum systems given by the Bures distance of the system state to the set of classical states with respect to one subsystem, that is, to the states with zero quantum discord. Our measure is a geometrical version of the quantum discord. As the latter it quantifies the degree of non-classicality in the system. For pure states it is identical to the geometric measure of entanglement. We show that for mixed states it coincides with the optimal success probability of an ambiguous quantum state discrimination task. Moreover, the closest zero-discord states to a state ρ are obtained in terms of the corresponding optimal measurements. By solving the quantum state discrimination problem for two qubits we determine the geometric quantum discord and the closest zero-discord states explicitly for states ρ with maximally disordered marginals.

PACS numbers: 03.67.Mn, 03.67.Hk

---

[*] Dominique.Spehner@ujf-grenoble.fr

I. INTRODUCTION

One of the basic question in quantum information theory is to understand how quantum correlations in composite quantum systems can be used to perform tasks that cannot be performed classically, or that lead classically to much lower classical efficiencies [1]. These correlations have been long thought to come solely from the entanglement among the different subsystems. This is the case for quantum computation and communication protocols using pure states. For composite systems in mixed states, however, there is now increasing evidence that other types of quantum correlations, captured by the quantum discord of Ollivier and Zurek [2] and Henderson and Vedral [3], could provide the main resource to exploit in order to outperform classical algorithms [4–6] or in some communication protocols [7–9]. The quantum discord quantifies the amount of mutual information not accessible by local measurement on one subsystem. One can generate mixed states with nonzero discord but no entanglement by preparing locally statistical mixtures of nonorthogonal states, which cannot be perfectly discriminated by measurements. The strongest hint suggesting that the quantum discord may in certain cases quantify the resource responsible of quantum speedups is provided by the deterministic quantum computation with one qubit (DQC1) of Knill and Laflamme [10] leading to exponential speedups with respect to known classical algorithms. Even though no entanglement is present during the whole computation [11], a nonzero quantum discord appears between the pure single qubit and the other qubits [4, 5]. This can be nicely interpreted by using the monogamy relation [12] between the quantum discord of a bipartite system AB and the entanglement of B with its environment E if ABE is in a pure state [6].

A mathematically appealing way to quantify quantum correlations in multipartite systems is given by the minimal distance of the system state to a separable state [13]. The Bures metric [14, 15] provides a nice distance d_B on the convex cone of density matrices, which has better properties than the Hilbert-Schmidt distance d_2 from a quantum information perspective. In particular, d_B is monotonous and riemannian [16] and its metric coincides with the quantum Fisher information [17] playing an important role in high precision interferometry. As a consequence, the minimal Bures distance to separable states satisfies all criteria of an entanglement measure [13], which is not the case for the distance d_2. This entanglement measure has been widely studied in the literature [18–20]. By analogy with entanglement, a geometric measure of quantum discord has been defined by Dakić, Vedral, and Brukner [21] as the minimal distance of the system state to the set of zero-discord states. This geometric quantum discord has been evaluated explicitly for two qubits [21]. However, the aforementioned authors use the Hilbert-Schmidt distance d_2.

The aim of this work is to study a similar geometric quantum discord (GQD) as in Ref. [21] but based on the Bures distance d_B, which seems a more natural choice. This distance measure of quantum correlations has moreover a clearer geometrical interpretation than other measures [13, 22, 23] based on the relative entropy, which is not a distance on the set of density matrices. We show that it shares many of the properties of the quantum discord. Most importantly, our geometrical approach provides further information not contained in the quantum discord itself. In fact, one can look for the closest state(s) with zero discord to a given state ρ, and hence learn something about the "position" of ρ with respect to the set of zero-discord states. We calculate the Bures-GQD explicitly for pure states of bipartite quantum systems and for mixed two-qubit states with maximally mixed marginals. In both situations we also determine the closest zero-discord state(s) and compare our results to those obtained by using the Hilbert-Schmidt distance. Our results are obtained by first establishing a link, which holds for arbitrary mixed states ρ in any bipartite system, between the Bures-GQD and the closest zero-discord states and the well-known minimal-error quantum state discrimination problem.

The task of discriminating states pertaining to a known set $\{\rho_1, \cdots, \rho_n\}$ of states ρ_i with prior probabilities η_i plays an important role in quantum communication and quantum cryptography. For instance, the set $\{\rho_1, \cdots, \rho_n\}$ can encode a message to be sent to a receiver. The sender chooses at random some states among the ρ_i's and gives them to the receiver, who is required to identify them and henceforth to decode the message. With this goal, the receiver performs a measurement on each state given to him by the sender. If the ρ_i are non-orthogonal, a perfect discrimination is impossible, so that the amount of sent information is smaller than in the case of orthogonal states. The best the receiver can do is to find the measurement that minimizes in some way his probability of equivocation. Two distinct strategies have been widely studied in the literature (see e.g. the review [24]). In the first one, the receiver seeks for a generalized measurement with $(n + 1)$ outcomes, allowing him to identify perfectly each state ρ_i but such that one of the outcomes leads to an inconclusive result (unambiguous quantum state discrimination). The probability of the inconclusive outcome must be minimized. In the second one, the receiver looks for a measurement with n outcomes yielding the maximal success probability $P_S = \sum_{i=1}^{n} \eta_i P_{i|i}$, where $P_{i|i}$ is the probability of the measurement outcome i given that the state is ρ_i. This strategy is called minimal error (or ambiguous) quantum state discrimination. The maximal success probability and optimal measurements are known explicitly for $n = 2$ [25], but no general solution has been found so far for more than two states (see, however, [26]) except when the ρ_i are related to each other by some symmetry and have equal probabilities η_i (see [24, 27] and references therein). However, several upper bounds on P_S^{opt} are known [28] and the discrimination task can be solved efficiently numerically [29]. Let us also stress that unambiguous and ambiguous discrimination have been implemented experimentally for pure states [30] and, more recently, for mixed states [31], by using polarized light.

Our main result shows that the Bures-GQD of a mixed state ρ is equal to the maximal success probability P_S in ambiguous state discrimination of a family of states ρ_i depending on ρ. Moreover, the closest zero-discord states to ρ are given in terms of the corresponding optimal von Neumann measurement.

The paper is organized as follows. The definitions of the quantum discords and the Bures distance are given in Sect. II, together with their main properties. In Sect. III we show that the Bures-GQD for a pure state coincides with the geometric measure of entanglement and is simply related to the highest Schmidt coefficient. We explain this fact by noting that the closest zero-discord states to a pure state are convex combinations of orthogonal product states. The link between the minimal Bures distance to the set of zero-discord states and ambiguous quantum state discrimination is explained and proved in the following Sect. IV. Finally, we study in Sec. V the case of two qubits. In that section the Bures-GQD is calculated explicitly and compared with the other quantum discords, and we determine the closest zero-discord states. The last section contains some conclusive remarks and perspectives.

II. DEFINITIONS OF THE QUANTUM DISCORDS

In this work we consider a bipartite quantum system AB with Hilbert space $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$, the spaces $\mathcal{H}_A$ and $\mathcal{H}_B$ of the subsystems A and B having arbitrary finite dimensions n_A and n_B. The states of AB are given by density matrices ρ on $\mathcal{H}$ (i.e., hermitian positive $N \times N$ matrices with unit trace, with $N = n_A n_B$). The reduced states of A and B are defined by partial tracing ρ over the other subsystem. They are denoted by $\rho_A = \operatorname{tr}_B(\rho)$ and $\rho_B = \operatorname{tr}_A(\rho)$.

Let us first recall the definition of the quantum discord [2, 3]. The total correlations of the bipartite system in the state ρ is described by the mutual information $I_{A:B}(\rho) = S(\rho_A) + S(\rho_B) - S(\rho)$, where $S(\cdot)$ stands for the von Neumann entropy. The amount $J_{B|A}(\rho)$ of classical correlations is given by the maximal reduction of entropy of the subsystem B after a von Neumann measurement on A. Such a measurement is described by an orthogonal family $\{\pi_i^A\}$ of projectors acting on $\mathcal{H}_A$. Hence $J_{B|A}(\rho) = \max_{\{\pi_i^A\}}\{S(\rho_B) - \sum_i q_i S(\rho_{B|i})\}$, where the maximum is over all von Neumann measurements $\{\pi_i^A\}$, $q_i = \operatorname{tr}(\pi_i^A \otimes 1\,\rho)$ is the probability of the measurement outcome i, and $\rho_{B|i} = q_i^{-1} \operatorname{tr}_A(\pi_i^A \otimes 1\,\rho)$ is the corresponding conditional state of B. The quantum discord is by definition the difference $\delta_A(\rho) = I_{A:B}(\rho) - J_{B|A}(\rho)$ between the total and classical correlations. It measures the amount of mutual information which is not accessible by local measurements on the subsystem A. Note that it is asymmetric under the exchange $A \leftrightarrow B$. It can be shown [32] that $\delta_A(\rho) \geq 0$ for any ρ, and $\delta_A(\sigma_{A\text{-cl}}) = 0$ if and only if

$$\sigma_{A\text{-cl}} = \sum_{i=1}^{n_A} q_i |\alpha_i\rangle\langle\alpha_i| \otimes \sigma_{B|i} \ , \tag{1}$$

where $\{|\alpha_i\rangle\}_{i=1}^{n_A}$ is an orthonormal basis of $\mathcal{H}_A$, $\sigma_{B|i}$ are some (arbitrary) states of B depending on the index i, and $q_i \geq 0$ are some probabilities, $\sum_i q_i = 1$. The fact that $\delta_A(\sigma_{A\text{-cl}}) = 0$ follows directly from $I_{A:B}(\sigma_{A\text{-cl}}) = S(\operatorname{tr}_A(\sigma_{A\text{-cl}})) - \sum_i q_i S(\sigma_{B|i}) \leq J_{B|A}(\sigma_{A\text{-cl}})$ and from the non-negativity of the quantum discord. For a bipartite system in the state $\sigma_{A\text{-cl}}$, the subsystem A is in one of the orthogonal states $|\alpha_i\rangle$ with probability q_i, hence A behaves as a classical system. For this reason, we will call *A-classical states* the zero-discord states of the form (1). We denote by $\mathcal{C}_A$ the set of all A-classical states. By using the spectral decompositions of the $\sigma_{B|i}$, any A-classical state $\sigma_{A\text{-cl}} \in \mathcal{C}_A$ can be decomposed as

$$\sigma_{A\text{-cl}} = \sum_{i=1}^{n_A} \sum_{j=1}^{n_B} q_{ij} |\alpha_i\rangle\langle\alpha_i| \otimes |\beta_{j|i}\rangle\langle\beta_{j|i}| \tag{2}$$

where, for any fixed i, $\{|\beta_{j|i}\rangle\}_{j=1}^{n_B}$ is an orthonormal basis of $\mathcal{H}_B$, and $q_{ij} \geq 0$, $\sum_{ij} q_{ij} = 1$ (note that the $|\beta_{j|i}\rangle$ need not be orthogonal for distinct i's). One defines similarly the set $\mathcal{C}_B$ of B-classical states, which are the states with zero quantum discord when the subsystem B is measured. A state which is both A- and B-classical possesses an eigenbasis $\{|\alpha_i\rangle \otimes |\beta_j\rangle\}_{i=1,j=1}^{n_A,n_B}$ of product vectors. It is fully classical, in the sense that a quantum system in this state can be "simulated" by a classical apparatus being in the state (i, j) with probability q_{ij}.

Let us point out that $\mathcal{C}_A$, $\mathcal{C}_B$, and the set of classical states $\mathcal{C}$ are not convex. Their convex hull is the set $\mathcal{S}$ of separable states. A state σ_{sep} is separable if it admits a convex decomposition $\sigma_{\text{sep}} = \sum_m q_m |\phi_m\rangle\langle\phi_m| \otimes |\psi_m\rangle\langle\psi_m|$, where $\{|\phi_m\rangle\}$ and $\{|\psi_m\rangle\}$ are (not necessarily orthogonal) families of unit vectors in $\mathcal{H}_A$ and $\mathcal{H}_B$ and $q_m \geq 0$, $\sum_m q_m = 1$. For pure states, A-classical, B-classical, classical and separable states all coincide. Actually, according to (2) the pure A-classical (and, similarly, the pure B-classical) states are product states.

The geometric quantum discord (GQD) of a state ρ of AB has been defined in [21] as the square distance of ρ to the set $\mathcal{C}_A$ of A-classical states, i.e.,

$$D_A^{(2)}(\rho) = d_2(\rho, \mathcal{C}_A)^2 \equiv \min_{\sigma_{A\text{-cl}} \in \mathcal{C}_A} d_2(\rho, \sigma_{A\text{-cl}})^2 \tag{3}$$

where $d_2(\rho,\sigma) = (\mathrm{tr}[(\rho-\sigma)^2])^{1/2}$ is the Hilbert-Schmidt distance. Instead of taking this distance, we use in this article the Bures distance

$$d_B(\rho,\sigma) = 2\big(1-\sqrt{F(\rho,\sigma)}\big) \tag{4}$$

where ρ and σ are two density matrices and $F(\rho,\sigma)$ is their fidelity [1, 14, 33],

$$F(\rho,\sigma) = \|\sqrt{\rho}\sqrt{\sigma}\|_1^2 = \big[\mathrm{tr}\big([\sqrt{\sigma}\rho\sqrt{\sigma}]^{1/2}\big)\big]^2 \,. \tag{5}$$

It is known that (4) defines a Riemannian distance on the convex cone $\mathcal{E} \subset \mathrm{Mat}(\mathbb{C}, \mathrm{N}\times\mathrm{N})$ of all states of AB. Its metric coincides (apart from a numerical factor) with the quantum Fisher information which plays an important role in quantum metrology [17]. Moreover, d_B satisfies the following properties [1, 33]: for any ρ, σ, σ_1, and $\sigma_2 \in \mathcal{E}$,
(i) convexity: if $\eta_1, \eta_2 \geq 0$ and $\eta_1 + \eta_2 = 1$, then $d_B(\rho, \eta_1\sigma_1 + \eta_2\sigma_2) \leq \eta_1 d_B(\rho,\sigma_1) + \eta_2 d(\rho,\sigma_2)$;
(ii) d_B is preserved by unitary conjugations: if U is a unitary operator on $\mathcal{H}$, then $d_B(U\rho U^\dagger, U\sigma U^\dagger) = d_B(\rho,\sigma)$;
(iii) d_B is monotonous under the action of completely positive trace preserving maps T from $\mathrm{Mat}(\mathbb{C}, \mathrm{N}\times\mathrm{N})$ into itself: for any such T, $d_B(T\rho, T\sigma) \leq d_B(\rho,\sigma)$.

Note that the Hilbert-Schmidt distance d_2 also satisfies (ii) but fails to satisfy (iii). The monotonous Riemannian distances on $\mathcal{E}$ have been classified by Petz [16].

We define the geometric quantum discord (GQD) as

$$D_A(\rho) = d_B(\rho, \mathcal{C}_A) = 2(1-\sqrt{F_A(\rho)}) \quad , \quad F_A(\rho) = \max_{\sigma_{A\text{-cl}} \in \mathcal{C}_A} F(\rho, \sigma_{A\text{-cl}}) \,. \tag{6}$$

The property (ii) above implies that D_A and $D_A^{(2)}$ are invariant under conjugations by local unitaries, $\rho \mapsto U_A \otimes U_B \rho U_A^\dagger \otimes U_B^\dagger$, since such transformations leave $\mathcal{C}_A$ invariant. By property (iii), D_A is monotonous under local operations involving von Neumann measurements on A, generalized measurements on B, and classical communication.

By analogy with (6) one can define two other geometrical measures of quantum correlations: the distance $D(\rho) = d_B(\rho, \mathcal{C})$ to the set of classical states $\mathcal{C}$ and the geometric measure of entanglement $E(\rho) = d_B(\rho, \mathcal{S}) = 2(1-\sqrt{F_{\mathcal{S}}(\rho)})$, where $F_{\mathcal{S}}(\rho)$ is the maximal fidelity between ρ and a separable state $\sigma_{\mathrm{sep}} \in \mathcal{S}$. The second measure satisfies all criteria of an entanglement measure [13] (in particular, it is monotonous under local operations and classical communication by the property (iii)). It is related to other entanglement measures defined via a convex roof construction thanks to the identity [18]

$$F_{\mathcal{S}}(\rho) = \max_{\{p_m\},\{|\Psi_m\rangle\}} \sum_m p_m F_{\mathcal{S}}(|\Psi_m\rangle) \,, \tag{7}$$

where the maximum is taken over all pure state decompositions $\rho = \sum_m p_m |\Psi_m\rangle\langle\Psi_m|$ of ρ (with $\|\Psi_m\| = 1$ and $p_m \geq 0$, $\sum p_m = 1$) and $F_{\mathcal{S}}(|\Psi_m\rangle) = \max_{\sigma\in\mathcal{S}} F(|\Psi_m\rangle\langle\Psi_m|, \sigma)$. The measure $E(\rho)$ is a geometrical analog of the entanglement of formation [34]. The latter is defined via a convex roof construction from the von Neumann entropy of the reduced state, $E_{\mathrm{EoF}}(\rho) = \min_{\{p_m\},\{|\Psi_m\rangle\}} \sum_m p_m E_{\mathrm{EoF}}(|\Psi_m\rangle)$ with $E_{\mathrm{EoF}}(|\Psi_m\rangle) = S(\mathrm{tr}_B\, |\Psi_m\rangle\langle\Psi_m|)$.

Since $\mathcal{C} \subset \mathcal{C}_A \subset \mathcal{S}$, the three distances are ordered as

$$E(\rho) \leq D_A(\rho) \leq D(\rho) \,. \tag{8}$$

This ordering of quantum correlations is a nice feature of the geometric measures. In contrast, the entanglement of formation $E_{\mathrm{EoF}}(\rho)$ can be larger or smaller than the quantum discord $\delta_A(\rho)$ [35, 36].

III. THE BURES GEOMETRIC QUANTUM DISCORD OF PURE STATES

We first restrict our attention to pure states, for which one can obtain a simple formula for D_A in terms of the Schmidt coefficients μ_i. We recall that any pure state $|\Psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ admits a Schmidt decomposition

$$|\Psi\rangle = \sum_{i=1}^n \sqrt{\mu_i} |\varphi_i\rangle \otimes |\chi_i\rangle \tag{9}$$

where $n = \min\{n_A, n_B\}$ and $\{|\varphi_i\rangle\}_{i=1}^{n_A}$ (respectively $\{|\chi_j\rangle\}_{j=1}^{n_B}$) is an orthonormal basis of $\mathcal{H}_A$ ($\mathcal{H}_B$). The decomposition (9) is unique, the μ_i and $|\varphi_i\rangle$ (respectively $|\chi_j\rangle$) being the eigenvalues and eigenvectors of the reduced state ρ_A of A (respectively ρ_B of B).

Theorem 1. *If* $\rho_\Psi = |\Psi\rangle\langle\Psi|$ *is a pure state, then*

$$D_A(\rho_\Psi) = D(\rho_\Psi) = E(\rho_\Psi) = 2(1 - \sqrt{\mu_{\max}}) \tag{10}$$

where $\mu_{\max}$ *is the largest Schmidt coefficient* μ_i*. If moreover a single coefficient* μ_i *is equal to* $\mu_{\max}$ *then the closest* A*-classical (respectively classical, separable) state to* ρ_Ψ *is the pure product state* $\sigma = |\varphi_{\max} \otimes \chi_{\max}\rangle\langle\varphi_{\max} \otimes \chi_{\max}|$*, where* $|\varphi_{\max}\rangle$ *and* $|\chi_{\max}\rangle$ *are the vectors corresponding to* $\mu_{\max}$ *in the decomposition (9). If* $r > 1$ *Schmidt coefficients* μ_i *are equal to* $\mu_{\max}$*, say* $\mu_{\max} = \mu_1 = \ldots = \mu_r > \mu_{r+1}, \ldots, \mu_n$*, then infinitely many* A*-classical (respectively classical, separable) states* σ *minimize the distance* $d_B(\rho_\Psi, \sigma)$*. These closest states* σ *are convex combinations of the orthogonal product states* $|\alpha_l \otimes \beta_l\rangle\langle\alpha_l \otimes \beta_l|$*,* $l = 1, \ldots, r$*, with* $|\alpha_l\rangle = \sum_{i=1}^r u_{il}|\varphi_i\rangle$ *and* $|\beta_l\rangle = \sum_{i=1}^r u_{il}^*|\chi_i\rangle$*, where* $(u_{il})_{i,l=1}^r$ *is an arbitrary* $r \times r$ *unitary matrix and* $|\varphi_i\rangle$ *and* $|\chi_i\rangle$ *are the vectors in the decomposition (9).*

The expression (10) of the geometric measure of entanglement $E(\rho_\Psi)$ is basically known in the literature, and the closest separable states to pure and mixed states have been investigated in [18]. By inspection of (9) and (10), $D_A(\rho_\Psi) = 0$ if and only if $|\Psi\rangle$ is a product state, in agreement with the fact that A-classical pure states are product states (the same holds for the other quantum correlation measures D and E). Moreover, from the inequality $\mu_{\max} \geq 1/n$ (following from $\sum_{i=1}^n \mu_i = 1$) one deduces that $D_A(\rho_\Psi) \leq 2(1 - 1/\sqrt{n})$. The maximal value of D_A is reached when $\mu_i = 1/n$ for any i, that is, for the maximally entangled states (recall that such states are the pure states with maximal entropy $S(\rho_A) = -\sum_{i=1}^n \mu_i \ln \mu_i = \ln(n)$). Remarkably, the maximally entangled states are the pure states having the largest set of closest separable states (this set is a $(n^2 + n - 1)$ real-parameter submanifold of $\mathcal{E}$). For instance, in the case of two qubits (i.e., for $n_A = n_B = 2$), the Bell states $|\Phi^\pm\rangle = (|00\rangle \pm |11\rangle)/\sqrt{2}$ admit as closest separable states the classical states

$$\sigma_\pm = \sum_{l=0,1} q_l |\alpha_l\rangle\langle\alpha_l| \otimes |\beta_l\rangle\langle\beta_l| \quad , \quad |\alpha_l\rangle = u_{0l}|0\rangle + u_{1l}|1\rangle \quad , \quad |\beta_l\rangle = u_{0l}^*|0\rangle \pm u_{1l}^*|1\rangle \tag{11}$$

with $u_{0l}^* u_{0m} + u_{1l}^* u_{1m} = \delta_{ml}$ and $q_l \geq 0$, $q_0 + q_1 = 1$. Interestingly, typical decoherence processes (such as pure phase dephasing) transform $\rho_{\Phi^\pm}$ into one of its closest separable state $(|00\rangle\langle 00| + |11\rangle\langle 11|)/2$ at times $t \gg t_{\rm dec}$, where $t_{\rm dec}$ is the decoherence time. Slower relaxation processes modifying the populations in the states $|00\rangle$ and $|11\rangle$ do not further increase the distance to the initial state $\rho_{\Phi^\pm}$. The situation is different for a partially entangled state $|\Psi\rangle = \sqrt{\mu_0}|00\rangle + \sqrt{\mu_1}|11\rangle$ with $\mu_1 > \mu_0$: then the closest separable state is the pure state $|11\rangle$, but $|\Psi\rangle$ evolves asymptotically to a statistical mixture of $|00\rangle$ and $|11\rangle$ when the qubits are coupled e.g. to thermal baths at positive temperatures.

Proof. For a pure state ρ_Ψ, the fidelity reads $F(\rho_\Psi, \sigma_{A\text{-cl}}) = \langle\Psi|\sigma_{A\text{-cl}}|\Psi\rangle$. Replacing $\sigma_{A\text{-cl}}$ in (6) by the right-hand side of (2) we get

$$F_A(\rho_\Psi) = \max_{\{|\alpha_i\rangle\},\{|\beta_{j|i}\rangle\},\{q_{ij}\}} \Big\{ \sum_{ij} q_{ij} |\langle\alpha_i \otimes \beta_{j|i}|\Psi\rangle|^2 \Big\} = \max_{\|\alpha\|=\|\beta\|=1} |\langle\alpha \otimes \beta|\Psi\rangle|^2 \tag{12}$$

where we have used $\sum_{ij} q_{ij} = 1$. Thanks to (9) and to the Cauchy-Schwarz inequality, for any normalized vectors $|\alpha\rangle \in \mathcal{H}_A$ and $|\beta\rangle \in \mathcal{H}_B$ one has

$$\begin{aligned} |\langle\alpha \otimes \beta|\Psi\rangle| &\leq \sum_i \sqrt{\mu_i} \big|\langle\alpha|\varphi_i\rangle\langle\beta|\chi_i\rangle\big| \leq \sqrt{\mu_{\max}} \sum_i \big|\langle\alpha|\varphi_i\rangle\langle\beta|\chi_i\rangle\big| \\ &\leq \sqrt{\mu_{\max}} \Big[\sum_i |\langle\alpha|\varphi_i\rangle|^2\Big]^{1/2} \Big[\sum_i |\langle\beta|\chi_i\rangle|^2\Big]^{1/2} = \sqrt{\mu_{\max}}\ . \end{aligned} \tag{13}$$

Let us first assume that $\mu_1 = \mu_{\max} > \mu_2, \ldots, \mu_n$. Then $|\langle\alpha \otimes \beta|\Psi\rangle| = \sqrt{\mu_{\max}}$ if and only if $|\alpha\rangle = |\varphi_1\rangle$ and $|\beta\rangle = |\chi_1\rangle$. Thus the maximal fidelity between ρ_Ψ and an A-classical state is simply given by the largest Schmidt coefficient $\mu_{\max}$ and (10) follows from (6). Moreover, the maximum in the second member of Eq.(12) is reached when a single q_{ij} is non-vanishing, say $q_{ij} = \delta_{i1}\delta_{j1}$, and $|\alpha_1\rangle = |\varphi_1\rangle$, $|\beta_{1|1}\rangle = |\chi_1\rangle$. This means that the closest A-classical state to ρ_Ψ is the pure product state $|\varphi_1 \otimes \chi_1\rangle\langle\varphi_1 \otimes \chi_1|$.

More generally, let $\mu_1 = \ldots = \mu_r = \mu_{\max} > \mu_{r+1}, \ldots, \mu_n$. In order that the first and second inequalities in the first line of (13) be equalities, $\arg(\langle\alpha|\varphi_i\rangle\langle\beta|\chi_i\rangle) = \theta$ must be independent of i and either $|\alpha\rangle$ belongs to $V_{\max} = \operatorname{span}\{|\varphi_1\rangle, \ldots, |\varphi_r\rangle\}$ or $|\beta\rangle$ belongs to $\operatorname{span}\{|\chi_1\rangle, \ldots, |\chi_r\rangle\}$. Furthermore, the Cauchy-Schwarz inequality in the second line is an equality if and only if $|\langle\alpha|\varphi_i\rangle| = |\langle\beta|\chi_i\rangle|$ for all i. This shows that from any orthonormal family $\{|\alpha_l\rangle\}_{l=1}^r$ of $V_{\max}$, one can construct r orthogonal vectors $|\alpha_l \otimes \beta_l\rangle$ satisfying $|\langle\alpha_l \otimes \beta_l|\Psi\rangle| = \sqrt{\mu_{\max}}$ for all $l = 1, \ldots, r$, with $\langle\beta|\chi_l\rangle = e^{\mathrm{i}\theta}\langle\varphi_l|\alpha\rangle$. The probabilities $\{q_{ij}\}$ maximizing the sum inside the brackets in (12) are given by $q_{ij} = q_i$ if

$i=j\leq r$ and zero otherwise, where $\{q_l\}_{l=1}^r$ is an arbitrary set of probabilities. The corresponding A-classical states with maximal fidelities $F(\rho_\Psi,\sigma)$ are the classical states $\sigma=\sum_{l=1}^r q_l|\alpha_l\otimes\beta_l\rangle\langle\alpha_l\otimes\beta_l|$.

The equality between the correlation measures D_A, D, and E is a consequence of the fact that the closest states to ρ_Ψ are classical states. Such an equality is reminiscent from the equality between the entanglement of formation $E_{\rm EoF}$ and the quantum discord δ_A for pure states. Let us notice that it does *not* hold for the Hilbert-Schmidt distance, for which the closest A-classical state to a pure state is in general a mixed state. Actually, one infers from the expression

$$d_2(\rho_\Psi,\sigma_{A\text{-cl}})^2=\operatorname{tr}[(|\Psi\rangle\langle\Psi|-\sigma_{A\text{-cl}})^2]=1-2F(\rho_\Psi,\sigma_{A\text{-cl}})+\operatorname{tr}(\sigma_{A\text{-cl}}^2) \tag{14}$$

that the closest A-classical state results from a competition between the maximization of the fidelity $F(\rho_\Psi,\sigma_{A\text{-cl}})$ and the minimization of the trace $\operatorname{tr}(\sigma_{A\text{-cl}}^2)$, which is maximum for pure states. For instance, one can show that the closest A-classical state to the Bell states $|\Phi^\pm\rangle$ are mixed two-qubit states, see Sec V D. The validity of theorem 1 is one of the major advantage of the Bures-GQD over the Hilbert-Schmidt-GQD.

IV. THE BURES GEOMETRIC QUANTUM DISCORD OF MIXED STATES

A. Link with minimal error quantum state discrimination

The determination of $D_A(\rho)$ is much more involved for mixed states ρ than for pure states. We show in this section that this problem is related to ambiguous quantum state discrimination (QSD). As it has been recalled in the introduction, in ambiguous QSD a state ρ_i drawn from a known family $\{\rho_i\}_{i=1}^{n_A}$ with prior probabilities $\{\eta_i\}_{i=1}^{n_A}$ is sent to a receiver. The task of the latter is to determine which state he has been given with a maximal probability of success. To do so, he performs a generalized measurement and concludes that the state is ρ_j when his measurement result is j. The generalized measurement is given by a family of positive operators $M_i\geq 0$ satisfying $\sum_i M_i=1$ (POVM). The probability to find the result j is $P_{j|i}=\operatorname{tr}(M_j\rho_i)$ if the system is in the state ρ_i. The maximal success probability of the receiver reads

$$P_S^{\rm opt}(\{\rho_i,\eta_i\})=\max_{\text{POVM }\{M_i\}}\sum_{i=1}^{n_A}\eta_i\operatorname{tr}(M_i\rho_i)\;. \tag{15}$$

Theorem 2. *Let ρ be a state of the bipartite system AB with Hilbert space $\mathcal{H}=\mathcal{H}_A\otimes\mathcal{H}_B$ and $\alpha=\{|\alpha_i\rangle\}_{i=1}^{n_A}$ be an orthonormal basis of $\mathcal{H}_A$. Consider the subset of A-classical states with a diagonal marginal in the basis α,*

$$\mathcal{C}_A(\alpha)=\big\{\sigma_{A\text{-}cl}\in\mathcal{C}_A\;;\;\alpha\textit{ is an eigenbasis of }\operatorname{tr}_B(\sigma_{A\text{-}cl})\big\}\,. \tag{16}$$

Then the maximal fidelity $F(\rho,\mathcal{C}_A(\alpha))=\max_{\sigma_{A\text{-}cl}\in\mathcal{C}_A(\alpha)}F(\rho,\sigma_{A\text{-}cl})$ of ρ to this subset is equal to

$$F(\rho,\mathcal{C}_A(\alpha))=P_S^{\text{opt v.N.}}(\{\rho_i,\eta_i\})\equiv\max_{\{\Pi_i\}}\sum_{i=1}^{n_A}\eta_i\operatorname{tr}(\Pi_i\rho_i) \tag{17}$$

where $P_S^{\text{opt v.N.}}(\{\rho_i,\eta_i\})$ is the maximal success probability over all von Neumann measurements given by orthogonal projectors Π_i of rank n_B (that is, self-adjoint operators on $\mathcal{H}$ satisfying $\Pi_i\Pi_j=\delta_{ij}\Pi_i$ and $\dim(\Pi_i\mathcal{H})=n_B$), and

$$\eta_i=\langle\alpha_i|\rho_A|\alpha_i\rangle\qquad,\qquad\rho_i=\eta_i^{-1}\sqrt{\rho}|\alpha_i\rangle\langle\alpha_i|\otimes 1\sqrt{\rho} \tag{18}$$

(if $\eta_i=0$ then ρ_i is not defined but does not contribute to the sum in (17)).

This theorem will be proven in Sec. IV B. Note that the ρ_i are quantum states of AB if $\eta_i>0$, because the right-hand side of the last identity in (18) is a nonnegative operator and η_i is chosen such that $\operatorname{tr}(\rho_i)=1$. Moreover, $\{\eta_i\}_{i=1}^{n_A}$ is a set of probabilities (since $\eta_i\geq 0$ and $\sum_i\eta_i=\operatorname{tr}(\rho)=1$) and $\{\rho_i,\eta_i\}_{i=1}^{n_A}$ defines a convex decomposition of ρ, $\rho=\sum_i\eta_i\rho_i$.

Let us assume that ρ is invertible. Then the application of a result by Eldar [37] shows that the POVM maximizing the success probability $P_S(\{\rho_i,\eta_i\})$ in (15) is a von Neumann measurement with projectors Π_i of rank n_B, i.e.,

$$F(\rho,\mathcal{C}_A(\alpha))=P_S^{\text{opt v.N.}}(\{\rho_i,\eta_i\})=P_S^{\rm opt}(\{\rho_i,\eta_i\})\quad,\quad\rho>0\,. \tag{19}$$

In fact, one may first notice that all matrices ρ_i have rank $r_i=n_B$ (in fact, ρ_i has the same rank as $\eta_i\rho^{-1/2}\rho_i=|\alpha_i\rangle\langle\alpha_i|\otimes 1\sqrt{\rho}$ and the latter matrix has rank n_B). Next, we argue that the ρ_i are linearly independent, in the sense that their eigenvectors $|\xi_{ij}\rangle$ form a linearly independent family $\{|\xi_{ij}\rangle\}_{i=1,\ldots,n_A}^{j=1,\ldots,n_B}$ of vectors in $\mathcal{H}$. Actually, a necessary and sufficient condition for $|\xi_{ij}\rangle$ to be an eigenvector of ρ_i with eigenvalue $\lambda_{ij}>0$ is $|\xi_{ij}\rangle = (\lambda_{ij}\eta_i)^{-1}\sqrt{\rho}|\alpha_i\rangle\otimes|\zeta_{ij}\rangle$, $|\zeta_{ij}\rangle\in\mathcal{H}_B$ being an eigenvector of $R_i=\langle\alpha_i|\rho|\alpha_i\rangle$ with eigenvalue $\lambda_{ij}\eta_i>0$. For any i, the hermitian invertible matrix R_i admits an orthonormal eigenbasis $\{|\zeta_{ij}\rangle\}_{j=1}^{n_B}$. Thanks to the invertibility of $\sqrt{\rho}$, $\{|\xi_{ij}\rangle\}_{i=1,\ldots,n_A}^{j=1,\ldots,n_B}$ is a basis of $\mathcal{H}$ and thus the states ρ_i are linearly independent. It is shown in [37] that for such a family of linearly independent states the second equality in (19) holds true.

The following result on the Bures-GQD of mixed states is a direct consequence of Theorem 2.

Theorem 3. *For any state ρ of the bipartite quantum system AB, the fidelity to the closest A-classical state is given by*

$$F_A(\rho) = \max_{\{|\alpha_i\rangle\}}\max_{\{\Pi_i\}}\sum_{i=1}^{n_A}\mathrm{tr}[\Pi_i\sqrt{\rho}|\alpha_i\rangle\langle\alpha_i|\otimes 1\sqrt{\rho}] \tag{20}$$

where the maxima are over all orthonormal basis $\{|\alpha_i\rangle\}$ of $\mathcal{H}_A$ and all families of orthogonal projectors $\{\Pi_i\}_{i=1}^{n_A}$ of $\mathcal{H}_A\otimes\mathcal{H}_B$ with rank n_B. Hence, using the notation of Theorem 2,

$$F_A(\rho) = \max_{\{|\alpha_i\rangle\}} P_S^{\mathrm{opt\,v.N.}}(\{\rho_i,\eta_i\})\,. \tag{21}$$

If $\rho>0$ then one can replace $P_S^{\mathrm{opt\,v.N.}}$ in (21) by the maximal success probability (15) over all POVMs.

It is noteworthy to observe that the basis vectors $|\alpha_i\rangle$ can be recovered from the states ρ_i and probabilities η_i by forming the square root measurement operators $M_i=\eta_i\rho^{-1/2}\rho_i\rho^{-1/2}$, with $\rho=\sum_i\eta_i\rho_i$ (we assume here $\rho>0$). Actually, such measurement operators are equal to the rank-n_B projectors $M_i=|\alpha_i\rangle\langle\alpha_i|\otimes 1$. By bounding from below $P_S^{\mathrm{opt\,v.N.}}(\{\rho_i,\eta_i\})$ by the success probability corresponding to $\Pi_i=M_i$, we obtain

$$F_A(\rho)\geq\max_{\{|\alpha_i\rangle\}}\sum_{i=1}^{n_A}\mathrm{tr}_B\left[\langle\alpha_i|\sqrt{\rho}|\alpha_i\rangle^2\right]\,. \tag{22}$$

The square root measurement plays an important role in the discrimination of almost orthogonal states [38] and of ensembles of states with certain symmetries [27].

To illustrate our result, let us study the ambiguous QSD task for some specific states ρ.

(i) If ρ is an A-classical state, i.e., if it admits the decomposition (1), then the basis $\{|\alpha_i\rangle\}$ maximizing the optimal success probability in (21) coincides with the basis appearing in this decomposition. With this choice, one obtains $\eta_i=q_i$ and $\rho_i=|\alpha_i\rangle\langle\alpha_i|\otimes\sigma_{B|i}$ for all i such that $q_i>0$. The states ρ_i are orthogonal and can thus be perfectly discriminated by von Neumann measurements, so that $F_A(\rho)=P_S^{\mathrm{opt\,v.N.}}(\{\rho_i,\eta_i\})=1$. Reciprocally, if $F_A(\rho)=1$ then $P_S^{\mathrm{opt\,v.N.}}(\rho)=1$ for some basis $\{|\alpha_i\rangle\}$ of $\mathcal{H}_A$ and the corresponding ρ_i must be orthogonal, that is, $\rho_i=\Pi_i\rho_i\Pi_i$ for some family $\{\Pi_i\}$ of orthogonal projectors with rank n_B. Hence $\rho=\sum_i\eta_i\rho_i=\sum_i\eta_i\Pi_i\rho_i\Pi_i$, $\sqrt{\rho}=\sum_i\Pi_i\sqrt{\rho}\,\Pi_i$, and (18) entails $\eta_i\rho_i=\eta_i\Pi_i\rho_i\Pi_i=\sqrt{\rho}|\alpha_i\rangle\langle\alpha_i|\otimes 1\sqrt{\rho}=\sqrt{\rho}\,\Pi_i|\alpha_i\rangle\langle\alpha_i|\otimes 1\,\Pi_i\sqrt{\rho}$, implying $\Pi_i=|\alpha_i\rangle\langle\alpha_i|\otimes 1$. Thus ρ is A-classical. This was of course to be expected since $D_A(\rho)=0$ if and only if ρ is A-classical, see Sec. II. Therefore, we can interpret our result (21) as follows: *the non-zero discord states ρ are such that the states (18) are non-orthogonal and thus cannot be perfectly discriminated for any orthonormal basis $\{|\alpha_i\rangle\}$ of $\mathcal{H}_A$.*

(ii) If $\rho=\rho_\Psi$ is a pure state, then all ρ_i with $\eta_i>0$ are identical and equal to ρ_Ψ, so that $P_S^{\mathrm{opt}}=P_S^{\mathrm{opt\,v.N.}}=\sup_{\{\Pi_i\}}\sum_i\eta_i\langle\Psi|\Pi_i|\Psi\rangle=\eta_{\max}$. One gets back the result $F_A(\rho_\Psi)=\mu_{\max}$ of Sec. III by optimization over the basis $\{|\alpha_i\rangle\}$.

(iii) Let us determine the states ρ having the highest possible GQD, i.e., the smallest possible fidelity $F_A(\rho)$.

Proposition. *If $n_A\leq n_B$, the smallest fidelity $F_A(\rho)$ for all states ρ of AB is equal to $1/n_A$. If $rn_A\leq n_B<(r+1)n_A$ with $r=1,2,\ldots$, the states ρ with $F_A(\rho)=1/n_A$ are convex combinations of the r maximally entangled pure states $|\Psi_k\rangle=n_A^{-1/2}\sum_{i=1}^{n_A}|\phi_i^{(k)}\rangle\otimes|\psi_i^{(k)}\rangle$, $k=1,\ldots,r$, with $\langle\phi_i^{(k)}|\phi_j^{(k)}\rangle=\delta_{ij}$ and $\langle\psi_i^{(k)}|\psi_j^{(l)}\rangle=\delta_{kl}\delta_{ij}$.*

We deduce from this result that the GQD $D_A(\rho)$ varies between 0 and $2-2/\sqrt{n_A}$ if $n_A\leq n_B$. By virtue of Theorem 1, the proposition above, and the inequality $E(\rho)\leq D_A(\rho)$, the geometric measure of entanglement $E(\rho)$ also varies between these two values. This means that *the most distant states from the set of A-classical states $\mathcal{C}_A$ are also the most distant from the set of separable states $\mathcal{S}$.* If $n_A\leq n_B<2n_A$, these most distant states are maximally entangled pure states.

Proof. The success probability $P_S^{\text{opt v.N.}}$ must be clearly larger than the highest prior probability $\eta_{\max} = \max_i\{\eta_i\}$. (A receiver would obtain $P_S = \eta_{\max}$ by simply guessing that his state is $\rho_{i_{\max}}$, with $\eta_{i_{\max}} = \eta_{\max}$; a better strategy is of course to perform the von Neumann measurement $\{\Pi_i\}$ such that $\Pi_{i_{\max}}$ projects on a n_B-dimensional subspace containing the range of $\rho_{i_{\max}}$; this range has a dimension $\leq n_B$ by a similar argument as in the discussion following Theorem 2). In view of (21) and by using $\eta_{\max} \geq 1/n_A$ (since $\sum_i \eta_i = 1$) we get

$$F_A(\rho) \geq \frac{1}{n_A} \tag{23}$$

for any mixed state ρ.
When $n_A \leq n_B$ the bound (23) is optimum, the value $1/n_A$ being reached for the maximally entangled pure states $\rho = \rho_\Psi$, see Sec. III. Thus $1/n_A$ is the smallest possible fidelity. Let ρ be a state having such a fidelity $F_A(\rho) = 1/n_A$. According to (21) and since it has been argued before that $P_S^{\text{opt v.N.}} \geq \eta_{\max} \geq 1/n_A$, $F_A(\rho) = 1/n_A$ means that $P_S^{\text{opt v.N.}}(\{\rho_i, \eta_i\}) = 1/n_A$ whatever the orthonormal basis $\{|\alpha_i\rangle\}$. It is intuitively clear that this can happen only if the receiver gets a collection of identical states ρ_i with equal prior probabilities $\eta_i = 1/n_A$. A rigorous proof of this fact is given in the Appendix. From (18) and $\rho = \sum \eta_i \rho_i$ we then obtain $\langle\alpha_i|\rho_A|\alpha_i\rangle = 1/n_A$ and $\rho_i = \rho$ for any $i = 1, \ldots, n_A$ and any orthonormal basis $\{|\alpha_i\rangle\}$. The first equality implies $\rho_A = 1/n_A$. By replacing the spectral decomposition $\rho = \sum_k p_k |\Psi_k\rangle\langle\Psi_k|$ of ρ into (18), the second equality yields $\operatorname{tr}_B(|\Psi_k\rangle\langle\Psi_l|) = n_A^{-1}\delta_{kl}$ for all k, l with $p_k p_l \neq 0$. Taking advantage of this identity for $k = l$, one finds that the eigenvectors $|\Psi_k\rangle$ of ρ with positive eigenvalues p_k have all their Schmidt coefficients equal to $1/n_A$, i.e., their Schmidt decompositions read $|\Psi_k\rangle = n_A^{-1/2} \sum_{i=1}^{n_A} |\phi_i^{(k)}\rangle \otimes |\psi_i^{(k)}\rangle$. Moreover, $\operatorname{tr}_B(|\Psi_k\rangle\langle\Psi_l|) = 0$ is equivalent to $\mathcal{V}_B^{(k)} \perp \mathcal{V}_B^{(l)}$ with $\mathcal{V}_B^{(k)} = \operatorname{span}\{|\psi_i^{(k)}\rangle; i = 1, \ldots, n_A\} \subset \mathcal{H}_B$. If $n_B < (r+1)n_A$ then at most r subspaces $\mathcal{V}_B^{(k)}$ may be pairwise orthogonal. Thus at most r eigenvalues p_k are nonzero.

Let us briefly discuss the case $n_A > n_B$. In that case the smallest value of the maximal fidelity $F_{\mathcal{S}}(\rho)$ to a separable state (see Sec. II) is equal to $1/n_B$ and $F_{\mathcal{S}}(\rho) = 1/n_B$ if and only if ρ is a pure product state. This is a consequence of (7) and of the bound $F_{\mathcal{S}}(\rho_\Psi) \geq 1/n_B$ for pure states ρ_Ψ. As a result, $E(\rho)$ varies between 0 and $2 - 2/\sqrt{n}$ with $n = \min\{n_A, n_B\}$. We could not establish an analog result for the GQD $D_A(\rho)$. The following reasoning shows that the bound (23) is not optimal, i.e., there are no states ρ with fidelities $F_A(\rho)$ equal or close to $1/n_A$ when $n_A > n_B$. Indeed, by generalizing the proof above, such states should be close to pure states. Since for $n_A > n_B$ pure states have fidelities larger than $1/n_B$ (see Sect. III), one may expect that a state close to a pure state has a fidelity close to $1/n_B$ or larger. This can be shown rigorously by invoking the inequality $F_A(\rho) \geq \|\rho\|/n_B$, where $\|\rho\| = \max\{p_k\}$ is the norm of ρ. This inequality is obtained by bounding $F_A(\rho)$ by the maximal fidelity to the set of pure product states,

$$F_A(\rho) \geq \max_{\|\alpha\|=\|\beta\|=1} \langle\alpha\otimes\beta|\rho|\alpha\otimes\beta\rangle \geq \max_k\{p_k \mu_{\max}^{(k)}\} \geq \frac{\max\{p_k\}}{n_B}\,, \tag{24}$$

where we have used in the second inequality the argument in the proof of Theorem 1 to express $\max_{\|\alpha\|=\|\beta\|=1} |\langle\alpha\otimes\beta|\Psi_k\rangle|^2$ as the largest Schmidt coefficient $\mu_{\max}^{(k)}$ of $|\Psi_k\rangle$. Note that all bounds in (24) are optimal for pure states only.

B. Derivation of the variational formula (20)

To prove Theorems 2 and 3, we start by evaluating the trace norm in (5) by means of the formula $\|T\|_1 = \max_U |\operatorname{tr}(UX)|$, the maximum being over all unitary operators on $\mathcal{H}$. Using also (2), one gets

$$\begin{aligned}
\sqrt{F(\rho, \sigma_{A\text{-cl}}} &= \max_U \bigl|\operatorname{tr}(U\sqrt{\rho}\sqrt{\sigma_{A\text{-cl}}})\bigr| \\
&= \max_U \Bigl|\sum_{i,j} \sqrt{q_{ij}}\, \operatorname{tr}(U\sqrt{\rho}\, |\alpha_i\rangle\langle\alpha_i| \otimes |\beta_{j|i}\rangle\langle\beta_{j|i}|)\Bigr| \\
&= \max_{\{|\Phi_{ij}\rangle\}} \Bigl|\sum_{i,j} \sqrt{q_{ij}} \langle\Phi_{ij}|\sqrt{\rho}|\alpha_i\otimes\beta_{j|i}\rangle\Bigr| \\
&= \max_{\{|\Phi_{ij}\rangle\}} \sum_{i,j} \sqrt{q_{ij}} \bigl|\langle\Phi_{ij}|\sqrt{\rho}|\alpha_i\otimes\beta_{j|i}\rangle\bigr|\,.
\end{aligned} \tag{25}$$

In the third line we have replaced the maximum over unitaries U by a maximum over all orthonormal basis $\{|\Phi_{ij}\rangle\}$ of $\mathcal{H}$ (with $|\Phi_{ij}\rangle = U^\dagger|\alpha_i\otimes\beta_{j|i}\rangle$). The last equality in (25) can be explained as follows. The expression in the last line is clearly larger than that of the third line; since for any i and j one can choose the phase factors of the vectors $|\Phi_{ij}\rangle$ in such a way that $\langle \Phi_{ij}|\sqrt{\rho}|\alpha_i \otimes \beta_{j|i}\rangle \geq 0$, the two expressions are in fact equal.

One has to maximize the last member of (25) over all orthonormal basis $\{|\beta_{j|i}\rangle\}$ of $\mathcal{H}_B$ and all set of probabilities $\{q_{ij}\}$. The maximum over the probabilities q_{ij} is easy to evaluate by using the Cauchy-Schwarz inequality and $\sum_{i,j} q_{ij} = 1$. It is reached for

$$q_{ij} = \frac{|\langle \Phi_{ij}|\sqrt{\rho}|\alpha_i \otimes \beta_{j|i}\rangle|^2}{\sum_{ij} |\langle \Phi_{ij}|\sqrt{\rho}|\alpha_i \otimes \beta_{j|i}\rangle|^2} \; . \tag{26}$$

We thus obtain

$$F(\rho, \mathcal{C}_A(\alpha)) = \max_{\{|\beta_{j|i}\rangle\}} \max_{\{q_{ij}\}} F(\rho, \sigma_{A\text{-cl}}) = \max_{\{|\beta_{j|i}\rangle\}} \max_{\{|\Phi_{ij}\rangle\}} \sum_{i,j} \bigl| \langle \Psi_{j|i}|\beta_{j|i}\rangle \bigr|^2 \tag{27}$$

where we have set $|\psi_{j|i}\rangle = \langle \alpha_i|\sqrt{\rho}|\Phi_{ij}\rangle \in \mathcal{H}_B$. We proceed to evaluate the maximum over $\{|\beta_{j|i}\rangle\}$ and $\{|\Phi_{ij}\rangle\}$. Let us fix i and consider the orthogonal family of projectors of $\mathcal{H}$ of rank n_B defined by

$$\Pi_i = \sum_j |\Phi_{ij}\rangle\langle\Phi_{ij}| \; . \tag{28}$$

By the Cauchy-Schwarz inequality, for any fixed i, one has

$$\max_{\{|\beta_{j|i}\rangle\}} \sum_j \bigl| \langle \psi_{j|i}|\beta_{j|i}\rangle \bigr|^2 \leq \sum_j \|\psi_{j|i}\|^2 = \operatorname{tr}\bigl[\Pi_i \sqrt{\rho}|\alpha_i\rangle\langle\alpha_i| \otimes 1\, \sqrt{\rho}\bigr] \; . \tag{29}$$

Note that (29) is an inequality if the vectors $|\psi_{j|i}\rangle$ are orthogonal for different j's. We now show that this is the case provided that the $|\Phi_{ij}\rangle$ are chosen appropriately. In fact, let us take an arbitrary orthonormal basis $\{|\Phi_{ij}\rangle\}$ of $\mathcal{H}$ and consider the hermitian $n_B \times n_B$ matrix $S^{(i)}$ with coefficients given by the scalar products $S^{(i)}_{jk} = \langle \psi_{j|i}|\psi_{k|i}\rangle$. One can find a unitary matrix $V^{(i)}$ such that $\widetilde{S}^{(i)} = (V^{(i)})^\dagger S^{(i)} V^{(i)}$ is diagonal and has nonzero diagonal elements in the first r_i raws, where r_i is the rank of $S^{(i)}$. Let $|\widetilde{\Phi}_{ij}\rangle = \sum_{l=1}^{n_B} V^{(i)}_{lj} |\Phi_{il}\rangle$. Then $\{|\widetilde{\Phi}_{ij}\rangle\}$ is an orthonormal basis of $\mathcal{H}$ and $\sum_j |\widetilde{\Phi}_{ij}\rangle\langle\widetilde{\Phi}_{ij}| = \Pi_i$. Moreover, the vectors $|\widetilde{\psi}_{j|i}\rangle = \langle\alpha_i|\sqrt{\rho}|\widetilde{\Phi}_{ij}\rangle = \sum_{l=1}^{n_B} V^{(i)}_{lj}|\psi_{l|i}\rangle$ form an orthogonal set $\{|\widetilde{\psi}_{j|i}\rangle\}_{j=1}^{r_i}$ and vanish for $j > r_i$. Therefore, for any fixed orthogonal family $\{\Pi_i\}_{i=1}^{n_A}$ of projectors of rank n_B, there exists an orthonormal basis $\{|\Phi_{ij}\rangle\}$ of $\mathcal{H}$ such that (28) holds and the inequality in (29) is an equality. Substituting this equality into (27), one finds

$$F(\rho, \mathcal{C}_A(\alpha)) = \max_{\{\Pi_i\}} \sum_{i,j} \|\widetilde{\psi}_{j|i}\|^2 = \max_{\{\Pi_i\}} \sum_i \operatorname{tr}[\Pi_i \sqrt{\rho}|\alpha_i\rangle\langle\alpha_i| \otimes 1\, \sqrt{\rho}] \; , \tag{30}$$

which yields the result (17). The formula (20) is obtained by maximization over the basis $\{|\alpha_i\rangle\}$.

C. Closest A-classical states

The proof of the previous subsection also gives an algorithm to find the closest A-classical state to a given mixed state ρ. To this end, one must find the orthonormal basis $\{|\alpha_i^{\text{opt}}\rangle\}_{i=1}^{n_A}$ of $\mathcal{H}_A$ maximizing $P_S^{\text{opt v.N.}}(\{\rho_i, \eta_i\})$ in (21) and the optimal von Neumann measurement $\{\Pi_i^{\text{opt}}\}_{i=1}^{n_A}$ yielding the minimal error in the discrimination of the states ρ_i^{opt} with probabilities η_i^{opt} which are associated to $\{|\alpha_i^{\text{opt}}\rangle\}$ by the formula (18).

Theorem 4. *The closest A-classical state to ρ is*

$$\sigma_\rho = \frac{1}{F_A(\rho)} \sum_{i=1}^{n_A} |\alpha_i^{\text{opt}}\rangle\langle\alpha_i^{\text{opt}}| \otimes \langle\alpha_i^{\text{opt}}|\sqrt{\rho}\, \Pi_i^{\text{opt}} \sqrt{\rho}|\alpha_i^{\text{opt}}\rangle_A \tag{31}$$

where $\{|\alpha_i^{\text{opt}}\rangle\}_{i=1}^{n_A}$ and $\{\Pi_i^{\text{opt}}\}_{i=1}^{n_A}$ are such that $F_A(\rho) = \sum_i \operatorname{tr}[\Pi_i^{\text{opt}} \sqrt{\rho}|\alpha_i^{\text{opt}}\rangle\langle\alpha_i^{\text{opt}}| \otimes 1\, \sqrt{\rho}]$, see Eq.(20).

Proof. This proof follows directly from that of the previous subsection. Actually, by using the expression (26) of the optimal probabilities q_{ij} and the fact that the Cauchy-Schwarz inequality (29) is an equality if and only if

$|\beta_{j|i}\rangle = |\psi_{j|i}\rangle / \|\psi_{j|i}\|$ when $\|\psi_{j|i}\| \neq 0$, we conclude that the closest A-classical state to ρ is given by Eq.(2) with $|\alpha_i\rangle = |\alpha_i^{\rm opt}\rangle$ and

$$q_i \sigma_{B|i} = \sum_{j=1}^{n_B} q_{ij} |\beta_{j|i}\rangle\langle\beta_{j|i}| = \frac{\sum_{j=1}^{n_B} |\widetilde{\psi}_{j|i}\rangle\langle\widetilde{\psi}_{j|i}|}{\sum_{i=1}^{n_A}\sum_{j=1}^{n_B} \|\widetilde{\psi}_{j|i}\|^2} \tag{32}$$

The denominator is equal to $F_A(\rho)$, see (30). The numerator is the same as the second factor in the right-hand side of (31) as by construction $|\widetilde{\psi}_{j|i}\rangle = \langle\alpha_i^{\rm opt}|\sqrt{\rho}|\widetilde{\Phi}_{ij}\rangle$ and $\sum_j |\widetilde{\Phi}_{ij}\rangle\langle\widetilde{\Phi}_{ij}| = \Pi_i^{\rm opt}$.

V. THE TWO-QUBIT CASE

A. Maximal success probability

If the subsystem A is a qubit, the set $\{\rho_i\}$ contains only $n_A = 2$ states. The minimal error and optimal measurements to discriminate the states ρ_i with prior probabilities η_i are then easy to determine [24]. In our case we must maximize the success probability over all von Neumann measurements given by orthogonal projectors Π_0 and Π_1 having the same rank n_B. One starts by writing the projector Π_1 as $1 - \Pi_0$ in the expression of the success probability,

$$P_S^{\{\Pi_i\}}(\{\rho_i,\eta_i\}) = \eta_0 \,{\rm tr}(\Pi_0\rho_0) + \eta_1 \,{\rm tr}((1-\Pi_0)\rho_1) = \frac{1}{2}\big(1 - {\rm tr}\,\Lambda\big) + {\rm tr}(\Pi_0\Lambda) \tag{33}$$

with $\Lambda = \eta_0\rho_0 - \eta_1\rho_1$. Thanks to the min-max theorem, the maximum of ${\rm tr}(\Pi_0\Lambda)$ over all projectors Π_0 of rank n_B is equal to the sum of the n_B highest eigenvalues $\lambda_1 \geq \cdots \geq \lambda_{n_B}$ of the hermitian matrix Λ, and the optimal projector $\Pi_0^{\rm opt}$ is the spectral projector associated to these highest eigenvalues. One has

$$P_S^{\rm opt\,v.N.}(\{\rho_i,\eta_i\}) = \max_{\{\Pi_i\}} P_S^{\{\Pi_i\}}(\{\rho_i,\eta_i\}) = \frac{1}{2}\big(1 - {\rm tr}\,\Lambda\big) + \sum_{l=1}^{n_B} \lambda_l\ . \tag{34}$$

For the states ρ_i and probabilities η_i defined by (18), $\Lambda = \sqrt{\rho}\,(|\alpha_0\rangle\langle\alpha_0| - |\alpha_1\rangle\langle\alpha_1|) \otimes 1\,\sqrt{\rho}$ where $\{|\alpha_i\rangle\}_{i=0}^1$ is an orthonormal basis of $\mathbb{C}^2$. For any such basis the operator inside the parenthesis in the last formula is equal to $\sigma_{\mathbf{u}} = \sum_{m=1}^3 u_m \sigma_m$ for some unit vector $\mathbf{u} \in \mathbb{R}^3$, where σ_1, σ_2, and σ_3 are the Pauli matrices. Reciprocally, one can associate to any unit vector $\mathbf{u} \in \mathbb{R}^3$ the eigenbasis $\{|\alpha_i\rangle\}_{i=0}^1$ of $\sigma_{\mathbf{u}}$. Hence, by substituting (34) into (21) we obtain

$$F_A(\rho) = \frac{1}{2}\max_{\|\mathbf{u}\|=1}\Big\{1 - {\rm tr}\,\Lambda(\mathbf{u}) + 2\sum_{l=1}^{n_B}\lambda_l(\mathbf{u})\Big\} \quad , \quad \Lambda(\mathbf{u}) = \sqrt{\rho}\,\sigma_{\mathbf{u}} \otimes 1\,\sqrt{\rho}\ , \tag{35}$$

where $\lambda_l(\mathbf{u})$ are the eigenvalues of $\Lambda(\mathbf{u})$ in decreasing order. The term $1 - {\rm tr}\,\Lambda(\mathbf{u})$ is equal to ${\rm tr}(\rho_A(1-\sigma_{\mathbf{u}})) = 2\langle\alpha_1|\rho_A|\alpha_1\rangle$. Note that $-\rho \leq \Lambda(\mathbf{u}) \leq \rho$, so that $\sum_{l=1}^{n_B}\lambda_l(\mathbf{u}) \leq \sum_{l=1}^{n_B} p_l$, where p_l are the eigenvalues of ρ in decreasing order. If $\rho > 0$ then $\Lambda(\mathbf{u})$ has n_B positive and n_B negative eigenvalues; in such a case (34) reduces to the well known expression [24] $P_S^{\rm opt\,v.N.}(\{\rho_i,\eta_i\}) = P_S^{\rm opt}(\{\rho_i,\eta_i\}) = (1 + {\rm tr}\,|\Lambda(\mathbf{u})|)/2$.

In order to evaluate $F_A(\rho)$ one needs to diagonalize the $2n_B \times 2n_B$ matrix $\Lambda(\mathbf{u})$ and to determine the maximum over the unit vectors $\mathbf{u}$ in (35). We do this in the next subsection for a special class of two-qubit states.

B. Bures-GQD for states with maximally disordered marginals

With the aim of comparing D_A with the other quantum discords, let us consider a simple family of two-qubit states for which all discords can be easily calculated: the states ρ with maximally disordered marginals $\rho_A = \rho_B = 1/2$. Any such state can be written up to a conjugation by a local unitary $U_A \otimes U_B$ (which leaves all discords unchanged) as [39]

$$\rho = \frac{1}{4}\Big(1 \otimes 1 + \sum_{m=1}^{3} c_m \sigma_m \otimes \sigma_m\Big) \tag{36}$$

where the real vector $\mathbf{c} = (c_1, c_2, c_3)$ belongs to the tetrahedron with vertices $F_\pm = (\pm 1, \mp 1, 1)$ and $G_\pm = (\pm 1, \pm 1, -1)$, that is,

$$p_0 = \frac{1}{4}(1 - c_1 - c_2 - c_3) \geq 0\ ,\ p_m = \frac{1}{4}(1 + c_1 + c_2 + c_3 - 2c_m) \geq 0\ ,\ m = 1, 2, 3\ . \tag{37}$$

We now calculate $D_A(\rho)$ for two qubits in the state (36). In the standard basis $\{|00\rangle, |10\rangle, |01\rangle, |11\rangle\}$, the matrices ρ and $\sqrt{\rho}$ are given by

$$\rho = \frac{1}{4}\begin{pmatrix} 1+c_3 & 0 & 0 & c_1-c_2 \\ 0 & 1-c_3 & c_1+c_2 & 0 \\ 0 & c_1+c_2 & 1-c_3 & 0 \\ c_1-c_2 & 0 & 0 & 1+c_3 \end{pmatrix} \quad , \quad \sqrt{\rho} = \begin{pmatrix} t_0+t_3 & 0 & 0 & t_1-t_2 \\ 0 & t_0-t_3 & t_1+t_2 & 0 \\ 0 & t_1+t_2 & t_0-t_3 & 0 \\ t_1-t_2 & 0 & 0 & t_0+t_3 \end{pmatrix} \tag{38}$$

where

$$4t_0 = \sqrt{p_0} + \sqrt{p_1} + \sqrt{p_2} + \sqrt{p_3} \quad , \quad 4t_1 = -\sqrt{p_0} - \sqrt{p_1} + \sqrt{p_2} + \sqrt{p_3} \;, \tag{39}$$

with similar formulas for t_2 and t_3 obtained by permutation of the indices $1, 2, 3$. The real parameters t_ν satisfy $4(t_0^2 + t_1^2 + t_2^2 + t_3^2) = \operatorname{tr}\rho = 1$. Let θ and ϕ be the angles defined by the vector $\mathbf{u} = (\sin\theta\cos\phi, \sin\theta\sin\phi, \cos\theta)$ on the unit sphere. In the standard basis the matrix $\Lambda(\mathbf{u}) = \sqrt{\rho}\,\sigma_{\mathbf{u}} \otimes 1\sqrt{\rho}$ reads

$$\Lambda(\mathbf{u}) = \frac{1}{4}\begin{pmatrix} (a_3+b_3)\cos\theta & \zeta_\phi^*\sin\theta & \xi_\phi^*\sin\theta & 0 \\ \zeta_\phi\sin\theta & (a_3-b_3)\cos\theta & 0 & \xi_\phi^*\sin\theta \\ \xi_\phi\sin\theta & 0 & (-a_3+b_3)\cos\theta & \zeta_\phi^*\sin\theta \\ 0 & \xi_\phi\sin\theta & \zeta_\phi\sin\theta & (-a_3-b_3)\cos\theta \end{pmatrix} \tag{40}$$

with $\xi_\phi = a_1\cos\phi + \mathrm{i}a_2\sin\phi$, $\zeta_\phi = b_1\cos\phi + \mathrm{i}b_2\sin\phi$, and

$$b_m = 8(t_0^2 + t_m^2) - 1 \quad , \quad a_1 = 8(t_0t_1 + t_2t_3) \quad , \quad a_2 = 8(t_0t_2 + t_1t_3) \quad , \quad a_3 = 8(t_0t_3 + t_1t_2) \;. \tag{41}$$

One finds

$$a_3 = 2\big(-\sqrt{p_0p_3} + \sqrt{p_1p_2}\big) \quad , \quad b_3 = 2\big(\sqrt{p_0p_3} + \sqrt{p_1p_2}\big) \;. \tag{42}$$

Similar expressions holds for the other coefficients a_1, a_2, b_1, and b_2 by permuting the indices $1, 2, 3$.

The eigenvalues of $\Lambda(\mathbf{u})$ come in opposite pairs $(\lambda_\pm(\mathbf{u}), -\lambda_\pm(\mathbf{u}))$ with

$$\lambda_\pm(\mathbf{u}) = \frac{1}{4}\left|\sqrt{b_3^2\cos^2\theta + |\zeta_\phi|^2\sin^2\theta} \pm \sqrt{a_3^2\cos^2\theta + |\xi_\phi|^2\sin^2\theta}\right| \;, \tag{43}$$

in agreement with $\operatorname{tr}\Lambda(\mathbf{u}) = 1 - 2\langle\alpha_1|\rho_A|\alpha_1\rangle = 0$. But $|a_m| \leq b_m$ by (42), thus the second square root is smaller than the first one. One deduces from (35) that

$$F_A(\rho) = \frac{1}{2}\left(1 + \max_{\theta,\phi}\sqrt{b_3^2\cos^2\theta + (b_1^2\cos^2\phi + b_2^2\sin^2\phi)\sin^2\theta}\right) \;. \tag{44}$$

If the b_m are distinct from each other, the maximum over the angles θ and ϕ is reached for $\cos^2\theta, \cos^2\phi \in \{0, 1\}$, that is, for $\mathbf{u} = \pm\mathbf{e}_x$, $\pm\mathbf{e}_y$, or $\pm\mathbf{e}_z$. If $b_1 = b_2 > b_3$, this maximum is reached for $\cos^2\theta = 0$ and $\cos\phi$ arbitrary. Thus

$$F_A(\rho) = \frac{1+b}{2} \quad \text{and} \quad D_A(\rho) = 2\Big(1 - \sqrt{\frac{1+b}{2}}\Big) \tag{45}$$

with

$$\begin{aligned} b = \max_{m=1,\ldots,3}\{b_m\} = \frac{1}{2}\max\Big\{ & \sqrt{(1+c_1)^2 - (c_2-c_3)^2} + \sqrt{(1-c_1)^2 - (c_2+c_3)^2}\,, \\ & \sqrt{(1+c_2)^2 - (c_1-c_3)^2} + \sqrt{(1-c_2)^2 - (c_1+c_3)^2}\,, \\ & \sqrt{(1+c_3)^2 - (c_1-c_2)^2} + \sqrt{(1-c_3)^2 - (c_1+c_2)^2}\Big\} \;. \end{aligned} \tag{46}$$

We notice that if $|c_m|$ is maximum for $m = m_{\max}$ then this is also true for b_m, i.e., the maximal number inside the brackets in (46) is the $(m_{\max})$th member. Actually, one can show from (42) that

$$c_m^2 - c_k^2 = b_m^2 - b_k^2 \quad , \quad m, k = 1, \ldots, 3 \;. \tag{47}$$



FIG. 1. (Color online) (a) Tetrahedron $\mathcal{T}$ with vertices $F_\pm = (\pm 1, \mp 1, 1)$ and $G_\pm = (\pm 1, \pm 1, -1)$. The vectors $\mathbf{c}$ inside $\mathcal{T}$ represent physical states $\rho(\mathbf{c})$ with maximally disordered marginals. The A-classical states in $\mathcal{T}$ are located on the segments $[IK]$, $[JL]$, and $[MN]$ (thick red dashed lines), with $I(1,0,0)$, $J(0,1,0)$, $K(-1,0,0)$, $L(0,-1,0)$, $M(0,0,1)$, and $N(0,0,-1)$. The shaded region delimited by the octahedron $IJKLMN$ corresponds to separable states. (b) Cut view of $\mathcal{T}$ in the plane $c_3 = 0$. The state $\rho(\frac{1}{4}, \frac{1}{2}, 0)$ has a unique closest A-classical state $\sigma_\rho(0, s_2, 0)$ with $s_2 = (4+\sqrt{35}-\sqrt{3})/(8+\sqrt{35}+\sqrt{3}) \simeq 0.523$ (see (57)). States on the broken lines are at equal Bures distance from the segments $[IK]$ and $[JL]$. They have infinitely many closest A-classical states given by (60). On these lines $D_A(\rho)$ is not differentiable. We show the two closest states of the form (36) to the state $\tau(-\frac{1}{4}, -\frac{1}{4}, 0)$ (they are obtained by taking $\phi = 0$ and $\pi/2$ in (60)): $\sigma_\tau(s_1, 0, 0)$ and $\sigma_\tau(0, s_1, 0)$ with $s_1 = -(2+\sqrt{6}-\sqrt{2})/(2+\sqrt{6}+\sqrt{2}) \simeq 0.518$. (c) Cut view of $\mathcal{T}$ in the plane $c_1 = c_2$. States in the shaded region are separable. The state $\rho(\frac{1}{2}, \frac{1}{2}, -\frac{3}{4})$ has a unique closest A-classical state $\sigma_\rho(0, 0, s_3)$, where $s_3 = -(5+\sqrt{33})/(9+\sqrt{33}) \simeq -0.729$. The states inside the triangles $H_\mp O G_\pm$ (here $O(0,0,0)$ is the origin) admit infinitely many closest A-classical states given by (60). States belonging to the blue square and triangle in panels (b) and (c) and to a face of $\mathcal{T}$ in panel (a) have infinitely many closest states given by Eqs.(54), (56), and (55).

It is clear on (46) that the vectors $\mathbf{c}$ such that b takes the highest possible value $b = 1$ have a single non-vanishing component c_m. Therefore, in agreement with Ref. [21], the A-classical states with maximally disordered marginals are situated (up to local unitary equivalence) on the three segments of the coordinate axes inside the tetrahedron, see Fig. 1(a). As a consequence, the separable states with maximally disordered marginals are located inside an octahedron with vertices on the three coordinate axes at a distance unity from the origin. This result is well-known [39]. It follows here immediately from the fact that such separable states are convex combinations of A-classical states. For indeed, the octahedron is the convex hull of the three aforementioned segments.

The states (36) with the highest GQD $D_A = 2 - \sqrt{2}$ have $b_1 = b_2 = b_3 = 0$. In agreement with the proposition of Sec. IV A, these states are the four Bell states $|\Phi^\pm\rangle = (|00\rangle \pm |11\rangle)/\sqrt{2}$ and $|\Psi^\pm\rangle = (|01\rangle \pm |10\rangle)/\sqrt{2}$ located at the vertices $F_\pm$ and $G_\pm$ of the tetrahedron (see Fig. 1(a)). Note that an analogous result holds for the Hilbert-Schmidt-GQD [21].

Let us remark that the inequality (22) is in fact an equality for two-qubit states ρ with maximally disordered marginals $\rho_A = \rho_B = 1/2$. Actually, for such states one has equal prior probabilities $\eta_i = \langle \alpha_i | \rho_A | \alpha_i \rangle = 1/2$ in the QSD problem of Theorem 2. Moreover, it follows from (36) that ρ is invariant under conjugation by the spin-flip operators $\sigma_1 \otimes \sigma_1$ and $\sigma_2 \otimes \sigma_2$. But it has been observed above that an optimal direction $\mathbf{u}^{\rm opt}$ is given by one of the coordinate vectors $\mathbf{e}_m$ or its opposite. Disregarding irrelevant phase factors, the two eigenvectors $|\alpha_0^{\rm opt}\rangle$ and $|\alpha_1^{\rm opt}\rangle$ of $\sigma_{\mathbf{u}^{\rm opt}}$ are transformed one into another by the spin-flip operator σ_m, with $m = 1$ if $\mathbf{u}^{\rm opt} = \pm\mathbf{e}_y, \pm\mathbf{e}_z$ and/or $m = 2$ if $\mathbf{u}^{\rm opt} = \pm\mathbf{e}_x, \pm\mathbf{e}_z$. Hence the states ρ_i are related by a unitary conjugation, $\rho_1 = \sigma_m \otimes \sigma_m \, \rho_0 \, \sigma_m \otimes \sigma_m$. The square root



FIG. 2. (Color online) Comparison of the various normalized quantum discords and geometric measure of entanglement for a Werner state $\rho_W = \frac{1-c}{4} 1 + c|\Psi^-\rangle\langle\Psi^-|$ with $|\Psi^-\rangle = (|01\rangle - |10\rangle)/\sqrt{2}$ (such a state is located on the segment $[OG_-]$ in Fig. 1(a)). From top to bottom: quantum discord δ_A (black boxes), Hilbert-Schmidt-GQD $\widetilde{D}_A^{(2)}$ (red crosses), Bures-GQD $\widetilde{D}_A$ (blue circles), and geometric measure of entanglement $\widetilde{E}$ (green diamonds).

measurement is known to be optimal to discriminate such symmetric set of states with equal prior probabilities [27]. Thus $F_A(\rho)$ is given by the right-hand side of Eq.(22).

C. Comparison with other quantum discords

We can now compare $D_A(\rho)$ for two-qubit states ρ of the form (36) with the other quantum discords defined in Sec. II. The corresponding quantum discord

$$\delta_A(\rho) = \sum_{\nu=0}^{3} p_\nu \ln_2 p_\nu + 2 - \frac{1-|c|}{2} \ln_2(1-|c|) - \frac{1+|c|}{2} \ln_2(1+|c|) \tag{48}$$

has been calculated in [40]. Here, $|c| = \max_m |c_m|$ and the probabilities p_ν are given by (37). The Hilbert-Schmidt-GQD defined by Eq.(3) is easy to determine for arbitrary two-qubit states [21]; for the particular states (36) it reads $D_A^{(2)}(\rho) = (\sum_m c_m^2 - |c|^2)/4$. Since the maximal values of D_A and $D_A^{(2)}$ are $2-\sqrt{2}$ and $1/2$, respectively, we renormalize the GQDs as

$$\widetilde{D}_A(\rho) = \frac{D_A(\rho)}{2-\sqrt{2}} \quad , \quad \widetilde{D}_A^{(2)}(\rho) = 2 D_A^{(2)}(\rho) \; . \tag{49}$$

We also compare $D_A(\rho)$ with the geometric measure of entanglement, given for two qubits by $E(\rho) = 2 - \sqrt{2}(1 + \sqrt{1 - C(\rho)^2})^{1/2}$ [18] where $C(\rho)$ is the Wootters concurrence. In our case $C(\rho) = 2\max\{|c_1 - c_2| - 1 + c_3, |c_1 + c_2| - 1 - c_3, 0\}$. The geometric measure of entanglement is normalized as $\widetilde{E}(\rho) = E(\rho)/(2-\sqrt{2})$.

In Fig. 2, the four quantum correlation measures $\widetilde{D}_A$, δ_A, $\widetilde{D}_A^{(2)}$, and $\widetilde{E}$ are plotted together for the Werner states, obtained by taking $c_1 = c_2 = c_3 = -c$ with $0 \leq c \leq 1$ in (36).

Fig. 3(a) displays the difference $\widetilde{D}_A(\rho) - \delta_A(\rho)$ for a family of states with maximally disordered marginals depending on two parameters. We observe on this figure that

$$\delta_A(\rho) \geq \widetilde{D}_A(\rho) \; . \tag{50}$$

In contrast, one sees in Figs. 2 and 3(b) that the Hilbert-Schmidt-GQD can be either smaller or larger than δ_A, as it has been noted previously in other works [41]. We have looked numerically for vectors **c** inside the tetrahedron violating (50) and have not found any such vector. This indicates that (50) is true for any state ρ with maximally disordered marginals. It would be of interest to know if it also holds true for more general two-qubit states.



FIG. 3. (Color online) (a) Difference $\delta_A(\rho) - \widetilde{D}_A(\rho)$ between the quantum discord and the normalized Bures-GQD for states with maximally disordered marginals with vectors $\mathbf{c}$ inside the triangle of vertices F_+, F_-, and N (see Fig. 1). The parameters u and v, $0 \leq u \leq 1 - v \leq 1$, are chosen such that $c_1 = -2u - v + 1 = -c_2$ and $c_3 = -2v + 1$. (b) same for the difference $\delta_A(\rho) - \widetilde{D}_A^{(2)}(\rho)$ between the quantum discord and the normalized Hilbert-Schmidt-GQD.

D. Closest A-classical states

We now turn to the problem of finding the A-classical two-qubit states σ_ρ satisfying

$$d_B(\rho, \sigma_\rho) = D_A(\rho) = \min_{\sigma_{A\text{-cl}} \in \mathcal{C}_A} d_B(\rho, \sigma_{A\text{-cl}}) \,. \tag{51}$$

These closest A-classical states to ρ provide useful information on the density matrix ρ and on the geometry of the set of quantum states. Such information is not contained in the discord $D_A(\rho)$ and can be of interest when studying dynamical evolutions, for instance, when the two qubits are coupled to their environments and undergo decoherence processes (see Sec. III). Like in the two previous subsections, we focus on states ρ with maximally disordered marginals given by Eq.(36), which form a 3-parameter subset $\mathcal{T}$ of the (15-parameter) set $\mathcal{E}$ of all two-qubit states. The geometric representation of the states $\rho \in \mathcal{T}$ by vectors $\mathbf{c}$ in the tetrahedron will give some insight to our results. Moreover, $\mathcal{T}$ contains many states of interest: maximally entangled states (the four Bell states on the vertices of the tetrahedron), the Werner states, separable states (located inside the octahedron), and A-classical states on the coordinate axes. The closest A-classical states for the Hilbert-Schmidt distance d_2 are also easy to determine for states $\rho \in \mathcal{T}$ by using the results of [21]. We will thus be able to compare the geometries on $\mathcal{T}$ associated to the two distances d_B and d_2.

According to Theorem 4, we must find the optimal basis $\{|\alpha_i^{\text{opt}}\rangle\}$ of $\mathcal{H}_A$ yielding the maximal value of the optimal success probability in (21) and the corresponding optimal measurement operators Π_0^{opt} and $\Pi_1^{\text{opt}} = 1 - \Pi_0^{\text{opt}}$. Changing the order of the vectors in the basis $\{|\alpha_i^{\text{opt}}\rangle\}$ amounts to exchange $(\rho_0^{\text{opt}}, \eta_0^{\text{opt}}) \leftrightarrow (\rho_1^{\text{opt}}, \eta_1^{\text{opt}})$ and $\Pi_0^{\text{opt}} \leftrightarrow \Pi_1^{\text{opt}}$. This does clearly not modify the optimal success probability and the A-classical state (31). But $\{|\alpha_i^{\text{opt}}\rangle\}$ is the eigenbasis of the spin matrix $\sigma_{\mathbf{u}}$ for the optimizing direction $\mathbf{u} = \mathbf{u}^{\text{opt}}$. Changing the order of the basis vectors thus corresponds to inverting $\mathbf{u}^{\text{opt}}$. Therefore, we need only to consider optimal directions $\mathbf{u}^{\text{opt}}$ satisfying $\mathbf{u}^{\text{opt}} \cdot \mathbf{e}_m \geq 0$ for $m = 1, 2$, or 3. Thanks to the results of Sec. V B, we know that if all components c_m of $\mathbf{c}$ have distinct moduli, then $\mathbf{u}^{\text{opt}} = \mathbf{e}_{m_{\max}}$, where $m_{\max}$ is the index m associated to the highest modulus $|c_m|$.

Let us first study the case $|c_3| > |c_1|, |c_2|$. Then $\mathbf{u}^{\text{opt}} = \mathbf{e}_3$ and $|\alpha_i^{\text{opt}}\rangle = |i\rangle$, $\{|i\rangle\}_{i=0}^{1}$ being the standard basis of $\mathcal{H}_A \simeq \mathbb{C}^2$. We have seen in Secs. V B and IV A that the states ρ_i^{opt} are optimally discriminated by the square root measurement given by the projectors $\Pi_i^{\text{opt}} = |\alpha_i^{\text{opt}}\rangle\langle\alpha_i^{\text{opt}}| \otimes 1$. In order to find all A-classical states σ_ρ satisfying (51) we must, however, find *all* von Neumann measurements $\{\Pi_i^{\text{opt}}\}$ maximizing the success probability $P_S(\{\rho_i^{\text{opt}}, \eta_i^{\text{opt}}\})$. As stressed in Sec. V A, Π_0^{opt} is the projector on the direct sum $\mathcal{V}_+ \oplus \mathcal{V}_-$ of the eigenspaces $\mathcal{V}_+$ and $\mathcal{V}_-$ associated to the eigenvalues $\lambda_+ \geq \lambda_- \geq 0$ of the matrix Λ, provided that $\lambda_\pm$ are non-degenerated. If $\lambda_+ > 0$ is simple and $\lambda_- = 0$ is two-fold degenerated, any projector on $\mathcal{V}_+ \oplus \mathcal{V}$ with $\mathcal{V} \subset \mathcal{V}_-$ a one-dimensional subspace of $\mathcal{V}_-$ defines an optimal projector $\Pi_0^{\rm opt}$. By setting $\theta = 0$ in (40) we find

$$\Pi_0^{\rm opt} = \begin{cases} |0\rangle\langle 0| \otimes 1 & \text{if } b_3 > |a_3| \\ |00\rangle\langle 00| + |\Phi\rangle\langle\Phi| & \text{if } b_3 = a_3 > 0, \quad \text{with } |\Phi\rangle \in \text{span}\{|10\rangle, |01\rangle\} \\ |01\rangle\langle 01| + |\Phi'\rangle\langle\Phi'| & \text{if } b_3 = -a_3 > 0, \text{ with } |\Phi'\rangle \in \text{span}\{|00\rangle, |11\rangle\}. \end{cases} \tag{52}$$

The condition $b_3 > |a_3|$ is achieved when all p_ν are nonzero, see (42). Then the optimal measurement $\{\Pi_i^{\rm opt}\}$ is unique and thus ρ has a unique closest A-classical state. The condition $b_3 = a_3$ (respectively $b_3 = -a_3$) corresponds to $p_0 p_3 = 0$ (respectively $p_1 p_2 = 0$), i.e., to a vector $\mathbf{c}$ belonging to one of the two faces $F_+F_-G_\pm$ (respectively $G_+G_-F_\pm$) of the tetrahedron. In this situation one has infinitely many measurements maximizing $P_S(\{\rho_i^{\rm opt}, \eta_i^{\rm opt}\})$. Note that (42) implies $b_3 \geq |a_3|$. Moreover, $p_0 p_3$ and $p_1 p_2$ cannot both vanish, since otherwise one would have $b_3 = a_3 = 0$, in contradiction with our hypothesis $|c_3| > |c_1|, |c_2|$ (which is equivalent to $b_3 > b_1, b_2 \geq 0$ by (47)). Replacing $|\alpha_i^{\rm opt}\rangle = |i\rangle$ and (52) into the expression (31) of σ_ρ, making use of the expression (38) of $\sqrt{\rho}$, and taking advantage of the identities $t_1 + t_2 = \pm(t_0 - t_3)$ for $p_{0,3} = 0$ and $t_1 - t_2 = \pm(t_0 + t_3)$ for $p_{1,2} = 0$ (which can be established with the help of (39)), one finds that σ_ρ is diagonal in the standard basis and has diagonal elements

$$\langle ii|\sigma_\rho|ii\rangle = \frac{(t_0 + t_3)^2}{F_A(\rho)}\big(1 + (-1)^i r'\big) \ , \ \langle ij|\sigma_\rho|ij\rangle = \frac{(t_0 - t_3)^2}{F_A(\rho)}\big(1 + (-1)^i r\big) \tag{53}$$

with $i, j = 0, 1$, $i \neq j$. Here, $r = \pm 2{\rm Re}\,\{\langle 01|\Phi\rangle\langle\Phi|10\rangle\}$ if $p_0 p_3 = 0$ and $r = 0$ otherwise, and $r' = \pm 2{\rm Re}\,\{\langle 11|\Phi'\rangle\langle\Phi'|00\rangle\}$ if $p_1 p_2 = 0$ and $r' = 0$ otherwise. If $p_0 p_3 = 0$ (respectively $p_1 p_2 = 0$) then r (respectively r') can take arbitrary values between -1 and 1, since $|\Phi\rangle$ ($|\Phi'\rangle$) is an arbitrary normalized vector in the two-dimensional space given in (52).

The case $|c_1| > |c_2|, |c_3|$ (respectively $|c_2| > |c_1|, |c_3|$) can be deduced from the previous case by the following symmetry argument. One can transform ρ into a state ρ' of the form (36) with a vector $\mathbf{c}' = (c_3, c_2, c_1)$ (respectively $\mathbf{c}' = (c_1, c_3, c_2)$) satisfying $|c_3'| > |c_1'|, |c_2'|$ by means of the unitary conjugation $\rho' = U\rho U^\dagger$ with $U = e^{{\rm i}\frac{\pi}{4}\sigma_2} \otimes e^{{\rm i}\frac{\pi}{4}\sigma_2}$ (respectively $U = e^{-{\rm i}\frac{\pi}{4}\sigma_1} \otimes e^{-{\rm i}\frac{\pi}{4}\sigma_1}$). By invariance of the Bures distance under unitary conjugations (property (ii) of Sec. II), such a transformation does not change the maximal fidelity, so that $F_A(\rho) = F_A(\rho') = (1 + b_3')/2 = (1 + b_{m_{\rm max}})/2$ with $m_{\rm max} = 1$ (respectively $m_{\rm max} = 2$), in agreement with (45). Moreover, according to (51) the states σ_ρ are related to the closest states $\sigma_{\rho'}$ to ρ' by $\sigma_\rho = U^\dagger \sigma_{\rho'} U$. Since $\sigma_{\rho'}$ has diagonal matrix elements in the standard basis, given by (53) upon the replacement of t_3 by $t_3' = t_{m_{\rm max}}$, this yields

$$\begin{aligned} \sigma_\rho = {} & \frac{(t_0 + t_{m_{\rm max}})^2}{F_A(\rho)}\Big((1 + r')|\alpha_0, \beta_0\rangle\langle\alpha_0, \beta_0| + (1 - r')|\alpha_1, \beta_1\rangle\langle\alpha_1, \beta_1|\Big) \\ & + \frac{(t_0 - t_{m_{\rm max}})^2}{F_A(\rho)}\Big((1 + r)|\alpha_0, \beta_1\rangle\langle\alpha_0, \beta_1| + (1 - r)|\alpha_1, \beta_0\rangle\langle\alpha_1, \beta_0|\Big) \end{aligned} \tag{54}$$

where we have set $|\alpha_i, \beta_j\rangle = |\alpha_i\rangle \otimes |\beta_j\rangle$ with $|\alpha_i\rangle = |\beta_i\rangle$ the eigenvectors of the Pauli matrix $\sigma_{m_{\rm max}}$, and

$$(r, r') \in \begin{cases} \{(0,0)\} & \text{if all } p_\nu \text{ are nonzero} \\ [-1, 1] \times \{0\} & \text{if } p_0 p_{m_{\rm max}} = 0 \text{ and } p_m > 0,\ m \neq m_{\rm max} \\ \{0\} \times [-1, 1] & \text{if } p_0 p_{m_{\rm max}} > 0 \text{ and } p_1 p_2 p_3 = 0. \end{cases} \tag{55}$$

The explicit expressions of the eigenvectors of $\sigma_{m_{\rm max}}$ in the standard basis read

$$|\alpha_i\rangle = |\beta_i\rangle = \begin{cases} |i\rangle & \text{if } |c_3| > |c_1|, |c_2| \\ 2^{-1/2}(|0\rangle + (-1)^i|1\rangle) & \text{if } |c_1| > |c_2|, |c_3| \\ 2^{-1/2}(|0\rangle + {\rm i}(-1)^i|1\rangle) & \text{if } |c_2| > |c_1|, |c_3|. \end{cases} \tag{56}$$

One checks on formulas (54) and (56) that when $\mathbf{c}$ does not pertain to one of the faces of the tetrahedron (so that $p_0 p_1 p_2 p_3 > 0$) and is such that $|c_m|$ is maximum for a single component $m = m_{\rm max}$, then σ_ρ is unique, has maximally disordered marginals, and is of the form

$$\sigma_\rho = \frac{1}{4}\Big(1 \otimes 1 + \frac{a_{m_{\rm max}} + c_{m_{\rm max}}}{2F_A(\rho)}\,\sigma_{m_{\rm max}} \otimes \sigma_{m_{\rm max}}\Big) \tag{57}$$

(we have used here the identities $F_A(\rho) = 4(t_0^2 + t_{m_{\rm max}}^2)$ and $16 t_0 t_m = a_m + c_m$, which follow from (37), (39), and (41)). The maximal fidelity $F_A(\rho)$ in the denominator in (57) is given by (45). Thanks to (42), one gets

$$a_1 = \frac{1}{2}\Big(\sqrt{(1 + c_1)^2 - (c_2 - c_3)^2} - \sqrt{(1 - c_1)^2 - (c_2 + c_3)^2}\Big) \tag{58}$$

with analogous formulas for a_2 and a_3 obtained via a permutation of indices.

We now turn to states ρ with vectors $\mathbf{c}$ such that $|c_m|$ is maximum for exactly two components c_m. For instance, let us assume $c_1 = \pm c_2$ and $|c_1| = |c_2| > |c_3|$ (the other cases will easily follow by a state rotation as above). Then $b_1 = b_2 > b_3$, see (47). Hence any vector $\mathbf{u}^{\rm opt}(\phi) = \cos\phi\,\mathbf{e}_x + \sin\phi\,\mathbf{e}_y$ in the (xOy)-plane defines an optimal direction (see Sec. V B). In other words, infinitely many basis $\{|\alpha_i^{\rm opt}\rangle\}$ maximize the optimal success probability. The situation is similar to the case of pure states with degenerate Schmidt coefficients discussed in Theorem 1. The non-uniqueness of $\mathbf{u}^{\rm opt}$ comes from the symmetry of ρ. For indeed, one finds from (36) that $\rho = U_\pm(\phi)\rho U_\pm(-\phi)$ for any ϕ, with $U_\pm(\phi) = e^{-{\rm i}\frac{\phi}{2}\sigma_3} \otimes e^{\mp{\rm i}\frac{\phi}{2}\sigma_3}$. By using the identity $e^{{\rm i}\frac{\phi}{2}\sigma_3}\sigma_{\mathbf{u}}e^{-{\rm i}\frac{\phi}{2}\sigma_3} = \sigma_{\mathbf{u}'}$ where $\mathbf{u}'$ is related to $\mathbf{u}$ by a rotation around the z-axis with the angle $-\phi$, it follows that

$$\Lambda\big(\mathbf{u}^{\rm opt}(\phi)\big) = U_\pm(\phi)\Lambda(\mathbf{e}_x)U_\pm(-\phi)\ . \tag{59}$$

As a result, the spectral projectors $\Pi_i^{\rm opt}(\phi)$ of $\Lambda\big(\mathbf{u}^{\rm opt}(\phi)\big)$ and thus the closest A-classical state (31) corresponding to $\mathbf{u}^{\rm opt}(\phi)$ are obtained from the corresponding projectors and A-classical states for $\mathbf{u}^{\rm opt} = \mathbf{e}_x$ by a unitary conjugation by $U_\pm(\phi)$. We conclude that all the A-classical states σ_ρ satisfying (51) are given

$$\sigma_\rho(\phi) = \frac{1}{4}\Big(1\otimes 1 + \frac{a_1+c_1}{2F_A(\rho)}\,(\cos\phi\,\sigma_1 + \sin\phi\,\sigma_2)\otimes(\cos\phi\,\sigma_1 \pm \sin\phi\,\sigma_2)\Big) \tag{60}$$

if $\mathbf{c}$ does not belong to a face of the tetrahedron. More generally, the closest states σ_ρ are of the form (54) with

$$|\alpha_i\rangle = \begin{cases} 2^{-\frac{1}{2}}(|0\rangle + (-1)^i e^{{\rm i}\phi}|1\rangle) \\ e^{-{\rm i}\frac{\theta}{2}\sigma_2}|i\rangle \\ e^{{\rm i}\frac{\theta}{2}\sigma_1}|i\rangle \end{cases} ,\ |\beta_i\rangle = \begin{cases} 2^{-\frac{1}{2}}(|0\rangle + (-1)^i e^{\pm{\rm i}\phi}|1\rangle) & \text{if } c_1 = \pm c_2,\ |c_1| > |c_3| \\ e^{\mp{\rm i}\frac{\theta}{2}\sigma_2}|i\rangle & \text{if } c_1 = \pm c_3,\ |c_1| > |c_2| \\ e^{\pm{\rm i}\frac{\theta}{2}\sigma_1}|i\rangle & \text{if } c_2 = \pm c_3,\ |c_2| > |c_1| \end{cases} \tag{61}$$

for arbitrary angles $0 \leq \phi < 2\pi$ and $0 \leq \theta < 2\pi$.

Finally, let us study the states ρ with $|c_1| = |c_2| = |c_3|$. This is the case for instance for the Werner states. Let $\epsilon_2, \epsilon_3 \in \{-1, 1\}$ be defined by $c_2 = \epsilon_2 c_1$ and $c_3 = \epsilon_3 c_1$. Then ρ is invariant under rotations with arbitrary angles θ and ϕ,

$$\rho = U_{\epsilon_2,\epsilon_3}(\theta,\phi)\rho\, U_{\epsilon_2,\epsilon_3}(\theta,\phi)^\dagger\ \ ,\ \ U_{\epsilon_2,\epsilon_3}(\theta,\phi) = e^{-{\rm i}\frac{\phi}{2}\sigma_3}e^{-{\rm i}\frac{\theta}{2}\sigma_2} \otimes e^{-{\rm i}\epsilon_2\frac{\phi}{2}\sigma_3}e^{-{\rm i}\epsilon_3\frac{\theta}{2}\sigma_2}\ . \tag{62}$$

Moreover, the eigenvalues of $\Lambda(\mathbf{u})$ are independent of $\mathbf{u}$, since $b_1 = b_2 = b_3$, see (43). Thus the optimal direction $\mathbf{u}^{\rm opt}$ is completely arbitrary and any orthonormal basis $\{|\alpha_i^{\rm opt}\rangle\}$ of $\mathbb{C}^2$ is optimal. The closest A-classical states to ρ are of the form (54) with

$$|\alpha_i\rangle = e^{-{\rm i}\frac{\phi}{2}\sigma_3}e^{-{\rm i}\frac{\theta}{2}\sigma_2}|i\rangle\ \ ,\ \ |\beta_i\rangle = e^{-{\rm i}\epsilon_2\frac{\phi}{2}\sigma_3}e^{-{\rm i}\epsilon_3\frac{\theta}{2}\sigma_2}|i\rangle\ . \tag{63}$$

The expressions in the standard basis of $|\alpha_i\rangle$ and $|\beta_i\rangle$ in (61) and (63) can be obtained from the relations

$$\begin{aligned} e^{-{\rm i}\epsilon_2\frac{\phi}{2}\sigma_3}e^{-{\rm i}\epsilon_3\frac{\theta}{2}\sigma_2}|0\rangle &= \cos\frac{\theta}{2}\,|0\rangle + \epsilon_3\,e^{{\rm i}\epsilon_2\phi}\sin\frac{\theta}{2}\,|1\rangle \\ e^{-{\rm i}\epsilon_2\frac{\phi}{2}\sigma_3}e^{-{\rm i}\epsilon_3\frac{\theta}{2}\sigma_2}|1\rangle &= \sin\frac{\theta}{2}\,|0\rangle - \epsilon_3 e^{{\rm i}\epsilon_2\phi}\cos\frac{\theta}{2}\,|1\rangle \end{aligned} \tag{64}$$

where we have disregarded irrelevant phase factors.

It should be noted that the closest A-classical states (54) are in fact classical states. This means that the distance of a state ρ with maximally disordered marginals to $\mathcal{C}_A$ coincides with its distance to the set $\mathcal{C}$ of classical states, i.e.,

$$D(\rho) = D_A(\rho) = D_B(\rho) = d_B(\rho,\sigma_\rho) = 2\Big(1 - \sqrt{\frac{1+b_{m_{\max}}}{2}}\Big)\ , \tag{65}$$

as could be expected from the symmetry of ρ under the exchange of the two qubits.

In Fig. 1 (b),(c), some examples of states ρ in the tetrahedron and their closest states σ_ρ are represented. Outside the dashed lines on which two or more components c_m have equal moduli, σ_ρ is specified by a vector $\mathbf{s}$ lying on the closest coordinate semi-axis to $\mathbf{c}$ for the usual distance in $\mathbb{R}^3$ (but $\mathbf{s}$ is not the closest vector to $\mathbf{c}$ on that semi-axis for this distance).

The closest A-classical states for the Hilbert-Schmidt distance d_2 to a state ρ of the form (36) can be found by using the results of Ref. [21]. They are given by

$$\sigma_\rho^{(2)} = \frac{1}{4}\Big(1\otimes 1 + \sum_{l,m=1}^{3} u_l c_m u_m \sigma_l \otimes \sigma_m\Big) \tag{66}$$

with an arbitrary unit vector $\mathbf{u} \in \mathbb{R}^3$. In contrast with the Bures distance, all states $\rho \in \mathcal{T}$ have infinitely many closest A-classical states $\sigma_\rho^{(2)}$. Moreover, there are three states $\sigma_\rho^{(2)}$ of the form (36) specified by vectors $\mathbf{s}_m = c_m \mathbf{e}_m$ lying on the coordinate axes $m = 1, 2, 3$, irrespective of the order of the moduli $|c_m|$. Since $\operatorname{tr}[(\sigma_\rho^{(2)})^2] = (1 + \sum_m c_m^2 u_m^2)/4 \leq 1/2$, the closest states $\sigma_\rho^{(2)}$ are always mixed. In particular, unlike for the distance d_B (see Theorem 1), the closest A-classical states to the maximally entangled pure states $|\Phi^\pm\rangle$ or $|\Psi^\pm\rangle$ are not pure product states.

VI. CONCLUSIONS

We have established in this paper a link between the ambiguous quantum state discrimination task and the problem of finding the minimal Bures distance of a state ρ to a state with zero quantum discord. More precisely, the maximal fidelity between ρ and an A-classical (i.e., zero-discord) state coincides with the maximal success probability in discriminating n_A states $\rho_i^{\rm opt}$ with prior probabilities $\eta_i^{\rm opt}$, given by Eq.(18) and depending on an optimal orthonormal basis $\{|\alpha_i^{\rm opt}\rangle\}_{i=1}^{n_A}$ of subsystem A (Theorem 3). The closest A-classical states to ρ are, in turn, given in terms of this optimal basis and of the optimal von Neumann measurements in the discrimination of $\{\rho_i^{\rm opt}, \eta_i^{\rm opt}\}$ (Theorem 4). By solving the quantum state discrimination problem we have found explicit formulas for the Bures-GQD and for the closest A-classical states to a given two-qubit state with maximally disordered marginals.

Since the quantum state discrimination task can be solved for $n_A = 2$ states, one can in principle by our method determine the GQD and the closest A-classical states for arbitrary two-qubit states, or even for arbitrary states of a bipartite system composed of a qubit A and a subsystem B with Hilbert space dimension $n_B > 2$. An open question for two qubits concerns the bound (50) between the quantum discord and the Bures-GQD, which holds for two-qubit states with maximally disordered marginals but could be more general. For subsystems A with higher Hilbert space dimensions $n_A > 2$, several open issues deserve further studies. Firstly, it would be desirable to characterize the "most quantum" states having the smallest fidelity to A-classical states when $n_A > n_B$. Secondly, it is not excluded that the specific quantum state discrimination task associated to the minimal Bures distance admits an explicit solution. Thirdly, the relation of the GQD with the geometric measure of entanglement should be investigated; in particular, there may exist some inequality analog to the monogamy relation [12] between the quantum discord and the entanglement of formation.

Our results on the closest A-classical states may shed new light on dissipative dynamical processes involving decoherence, i.e., evolutions towards classical states. In fact, our analysis allows to determine the geodesics passing through a given state ρ_0 with a non-zero discord. Such a geodesic contains all states ρ having the same closest A-classical state σ_{ρ_0} as ρ_0. It would be of interest to compare these Bures geodesics with the actual paths followed by the density matrix in specific physical examples.

Acknowledgements: We are grateful to Maria de los Angeles Gallego for revising the derivation of Eq.(20) and for providing us some notes about the calculation of the geometric discord for Werner states. D.S. thanks Aldo Delgado for interesting discussions. We acknowledge financial support from the french project ANR-09-BLAN-0098-01, the Chilean Fondecyt project no.100039, and the project Conicyt-PIA anillo ACT-1112 "Red de Análisis estocástico y aplicaciones".

Appendix A: A necessary and sufficient condition for the optimal success probability to be equal to the inverse number of states

Let $\{\rho_i\}_{i=1}^{n_A}$ be a family of n_A states on $\mathcal{H}$ with prior probabilities η_i, where $n_A = N/n_B$ is a divisor of $\dim(\mathcal{H}) = N$. We assume that the ρ_i have ranks $\operatorname{rank}(\rho_i) \leq n_B$ for any i. Let $P_S^{\rm opt\,v.N.}(\{\rho_i, \eta_i\})$ be the optimal success probability defined by Eq.(17). We prove in this appendix that $P_S^{\rm opt\,v.N.}(\{\rho_i, \eta_i\}) = 1/n_A$ if and only if $\eta_i = 1/n_A$ for any i and all states ρ_i are identical.

The conditions $\eta_i = 1/n_A$ and $\rho_i = \rho$ are clearly sufficient to have $P_S^{\rm opt\,v.N.}(\{\rho_i, \eta_i\}) = 1/n_A$ (a measurement cannot distinguish the identical states ρ_i and thus cannot do better than a random choice with equal probabilities). We need to show that they are also necessary conditions. Let us assume $P_S^{\rm opt\,v.N.}(\{\rho_i, \eta_i\}) = 1/n_A$. The equality of the prior probabilities η_i is obvious from the bounds $P_S^{\rm opt\,v.N.}(\{\rho_i, \eta_i\}) \geq \eta_{\max}$ and $\eta_{\max} \geq 1/n_A$ (see Sec. IV A, which implies $\eta_{\max} = 1/n_A = \eta_i$ for all i. As a result and by using our hypothesis, any orthogonal family $\{\Pi_i\}_{i=1}^{n_B}$ of projectors of rank n_B satisfies $\sum_i \operatorname{tr}(\Pi_i \rho_i) = n_A P_S(\{\rho_i, \eta_i\}) = 1$. We now argue that the states ρ_i have ranges contained in a common n_B-dimensional subspace. In fact, let $\mathcal{V}_1$ be the n_B-dimensional subspace of $\mathcal{H}$ containing the eigenvectors of ρ_1 associated to the n_B highest eigenvalues, and let us denote by Π_1 the projector onto $\mathcal{V}_1$. Then $\operatorname{ran} \rho_1 \subset \mathcal{V}_1$ (since we have assumed $\mathrm{rank}(\rho_1) \leq n_B$) and thus $\rho_1 = \Pi_1 \rho_1$. This implies $1 = \sum_i \mathrm{tr}(\Pi_i \rho_i) \geq \mathrm{tr}(\Pi_1 \rho_1) = 1$. It follows that $\mathrm{tr}(\Pi_2 \rho_2) = 0$ for any projector Π_2 of rank n_B orthogonal to Π_1. Hence ρ_1 and ρ_2, and similarly all ρ_i, $i = 3, \ldots, n_A$, have ranges contained in $\mathcal{V}_1$. In order to show that all the states ρ_i are equal, we further consider, for each $1 \leq k \leq n_B$, some n_B-dimensional subspace $\mathcal{V}_1^{(k)}$ containing the eigenvectors associated to the k highest eigenvalues $\lambda_1, \ldots, \lambda_k$ of ρ_1, the other eigenvectors being orthogonal to $\mathcal{V}_1^{(k)}$ (then $\mathcal{V}_1^{(n_B)} = \mathcal{V}_1$). We also choose a n_B-dimensional subspace $\mathcal{W}_1^{(k)} \subset \mathcal{H}$ orthogonal to $\mathcal{V}^{(k)}$ such that $\mathcal{W}_1^{(k)} \oplus \mathcal{V}_1^{(k)} \supset \mathcal{V}_1$. Let $\{\Pi_i^{(k)}\}_{i=1}^{n_A}$ be an orthogonal family of projectors of rank n_B such that $\Pi_1^{(k)}$ and $\Pi_2^{(k)}$ are the projectors onto $\mathcal{V}_1^{(k)}$ and $\mathcal{W}_1^{(k)}$, respectively. Then

$$1 = \sum_i \mathrm{tr}(\Pi_i^{(k)} \rho_i) = \mathrm{tr}(\Pi_1^{(k)} \rho_1) + \mathrm{tr}[(1 - \Pi_1^{(k)}) \rho_2] = 1 + \lambda_1 + \ldots + \lambda_k - \mathrm{tr}(\Pi_1^{(k)} \rho_2) \tag{A1}$$

where we used $\sum_i \Pi_i^{(k)} = 1$ and $\mathrm{ran}(\rho_i) \subset \mathcal{W}_1^{(k)} \oplus \mathcal{V}_1^{(k)}$ in the second equality. By virtue of the min-max theorem, $\mathrm{tr}(\Pi_1^{(k)} \rho_2)$ is smaller than the sum of the k highest eigenvalues of ρ_2. The identities (A1) imply that this sum is larger than the sum $\lambda_1 + \ldots + \lambda_k$ of the k highest eigenvalues of ρ_1. By exchanging the roles of ρ_1 and ρ_2 we obtain the reverse equality. Since moreover k is arbitrary between 1 and n_B, it follows that ρ_1 and ρ_2 have identical eigenvalues. By using (A1) again, $\mathrm{tr}(\Pi_1^{(k)} \rho_2)$ is equal to the sum of the k highest eigenvalues of ρ_2, implying that the k corresponding eigenvectors of ρ_2 are contained in $\mathcal{V}_1^{(k)}$. Since k is arbitrary, this proves that ρ_1 and ρ_2 have identical eigenspaces. Therefore $\rho_1 = \rho_2$. Repeating the same argument for the other states ρ_i, $i \geq 3$, we obtain $\rho_1 = \ldots = \rho_{n_A}$.

---

[1] N.A. Nielsen and I.L. Chuang, *Quantum Computation and Information* (Cambridge University Press, 2000)
[2] H. Ollivier and W.H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001)
[3] L. Henderson and V. Vedral, J. Phys. A **34**, 6899 (2001)
[4] A. Datta, A. Shaji, and C. Caves, Phys. Rev. Lett **100**, 050502 (2008)
[5] B.P. Lanyon, M. Barbieri, M.P. Almeida, and A.G. White, Phys. Rev. Lett. **101**, 200501 (2008)
[6] F.F. Fanchini, M.F. Cornelio, M.C. de Oliveira, and A.O. Caldeira, Phys. Rev. A **84**, 012313 (2011)
[7] V. Madhok and A. Datta, Phys. Rev. A **83**, 032323 (2011)
[8] M. Gu, H. M. Chrzanowski, S. M. Assad, T. Symul, K. Modi, T. C. Ralph, V. Vedral, and P. Koy Lam, Nature Physics **8**, 671 (2012)
[9] B. Dakić *et al.*, Nature Physics **8**, 666 (2012)
[10] E. Knill and R. Laflamme, Phys. Rev. Lett. **81**, 5672 (1998)
[11] A. Datta, S. T. Flammia, and C. M. Caves, Phys. Rev. A **72**, 042316 (2005)
[12] M. Koashi and A. Winter, Phys. Rev. A **69**, 022309 (2004)
[13] V. Vedral and M.B. Plenio, Phys. Rev. A **57**, 1619 (1998)
[14] A. Uhlmann, Rep. Math. Phys. **9** 273 (1976)
[15] D. Bures, Trans. Am. Math. Soc. **135**, 199 (1969)
[16] D. Petz, Lin. Alg. and its Appl. **244**, 81 (1996)
[17] S.L. Braunstein and C.M. Caves, Phys. Rev. Lett **72**, 3439 (1994)
[18] A. Streltsov, H. Kampermann, and D. Bruß, New. J. Phys. **12**, 123004 (2010)
[19] T.C. Wei and P.M. Goldbart, Phys. Rev. A **68**, 042307 (2003)
[20] L. Rudnicki, Z. Puchala, P. Horodecki, and K. Życzkowski, arXiv:1211.2306 [quant-ph]
[21] B. Dakić, V. Vedral, and C. Brukner, Phys. Rev. Lett **105**, 190502 (2010)
[22] K. Modi, T. Parerek, W. Son, V. Vedral, and M. Williamson, Phys. Rev. Lett. **104**, 080501 (2010)
[23] S. Friedland and G. Gour, J. Math. Phys. **52**, 052201 (2011)
[24] J. Bergou, U. Herzog, and M. Hillery, in: *Quantum state estimation*, Lecture Notes in Physics **649**, Eds. M. Paris and J. Rehacek (Springer, Berlin, 2004), pp. 417-465
[25] C.W. Helstrom, *Quantum Detection and Estimation Theory* (Academic Press, New York, 1976)
[26] J. Bae, arXiv:quant-phys/1210.2845
[27] Y.C. Eldar, A. Megretski, and G.C. Verghese, IEEE Trans. Inf. Theory **50**, 1198 (2004); C.-L. Chou and L.Y. Hsu, Phys. Rev. A **68**, 042305 (2003)
[28] D. Qiu and L. Li, Phys. Rev. A **81**, 042329 (2010)
[29] M. Jezek, J. Rehácek, and J. Fiurásek, arXiv:quant-ph/0201108
[30] B. Huttner *et al.*, Phys. Rev. A **54**, 3783 (1996); S.M. Barnett and E. Riis, J. Mod. Opt. **44**, 1061 (1997); R.B.M. Clarke *et al.*, Phys. Rev. A **63**, 040305(R) (2001); R.B.M. Clarke, V.M. Kendon, A. Chefles, S.M. Barnett, E. Riis, and M. Sasaki, Phys. Rev. A **64**, 012303 (2001)
[31] M. Mohseni, A.M. Steinberg, and J.A. Bergou, Phys. Rev. Lett. **93**, 200403 (2004)
[32] A. Datta, arXiv:quant-ph/1003.5256
[33] R. Jozsa, J. Mod. Opt. **94**, 2315 (1994)

[34] C.H. Bennett, D.P. DiVincenzo, J.A. Smolin, and W.K. Wootters, Phys. Rev. A **54**, 3824 (1996)
[35] M. de los Angeles Gallego and M. Orszag, J. Opt. Soc. Am. B **29**, 1690 (2012)
[36] M. de los Angeles Gallego, R. Coto, and M. Orszag, Phys. Scr. T **147**, 014012 (2012)
[37] Y.C. Eldar, Phys. Rev. A **68**, 052303 (2003)
[38] P. Hausladen and W.K. Wootters, J. Mod. Opt. **41**, 2385 (1994)
[39] R. Horodecki and M. Horodecki, Phys. Rev. A **54**, 1838 (1996)
[40] S. Luo, Phys. Rev. A **77**, 042303 (2008)
[41] D. Girolami and G. Adesso, Phys. Rev. A **83**, 052108 (2011)